Index

| absorption band 150, 168, 174, 176, 181 absorption peak 47, 154–157, 169, 171, 175, 176, 200, 207 absorption spectrum 152–154, 156, 157, 169, 172, 174–177, 181, 182, 200, 207 active site 232, 237, 240, 246, 247, 250, 257, 261, 265, 266, 271, 287, 308, 311–313 adsorption 203, 229, 234, 260, 263, 276, 300, 303, 304, 310, 312 asymmetric 308 bridging 246 dative amine 303 aggregation 69, 70, 186, 188, 266, 311 cation-induced 188 electron-rich induction 191 photo-induced 202 polymer-induced 191 solvent-induced 188, 191 aggregation-induced emission (AIE) 181, 185–188, 191, 192 AIE see aggregation-induced emission generic 303 primary 267, 268, 314 annealing 99, 231, 247 anthracene 234, 236 AuNC see gold nanocluster AuNP see gold nanoparticle | basin hopping (BH) 99, 100, 109 BE see binding energy BH see basin hopping binding energy (BE) 105, 125, |
|---|--|
| bandgap 156, 158, 207, 230 | polydisperse 230 |

| catalytic activity 230, 232, 235, | electronic transition 4, 153, 173, |
|-------------------------------------|------------------------------------|
| 237, 238, 243, 245, 257, 259, | 176, 231 |
| 269, 273–275, 277, 279, 280, | electron paramagnetic resonance |
| 314 | (EPR) 258, 294 |
| catalytic hydrogenation 232, 266, | electron transfer 125, 132, 181, |
| 268 | 198, 199, 201, 236, 259, 275, |
| catalytic mechanism 231, 250, | 295 |
| 256, 285, 293, 297, 306, 313, | emission 165, 191, 233 |
| 317, 318 | near-infrared 182 |
| catalytic oxidation 236, 240, 262 | optical 165 |
| catalytic performance 229, 230, | two-photon excitation 193 |
| 234, 235, 237, 246, 247, 256, | emission spectrum 184, 189, 194, |
| 266, 276, 277, 280, 297, 299, | 293, 294 |
| 303, 304 | enantiomer 22, 26, 27, 50, 79, 202 |
| catalytic properties 126, 255–257, | 205, 206, 208 |
| 266, 274, 280, 285, 287, 298 | energy 101, 152, 155, 172, 190, |
| catalytic reaction 229, 233, 235, | 195, 199, 269, 286, 301–303, |
| 243, 271, 273, 287, 288, 292, | 313 |
| 295-297, 300, 302, 314, | catalytic reaction 290 |
| 316-318 | cohesive 42 |
| catalytic reactivity 45, 70, 303, | dissociation 124 |
| 312, 317 | exchange-correlation 290 |
| chemical system 285, 290, 291, | hydrogenation reaction 308 |
| 317 | ionization 124 |
| chemoselective hydrogenation | molecular orbital 77 |
| 232, 244, 265 | photon 155 |
| chemoselectivity 232, 265 | pump pulse 155, 158, 160 |
| chiral clusters 197, 203, 204, 206 | renewable 272 |
| chirality 49, 51, 57, 58, 166, 197, | solar 295 |
| 202-205, 208 | thermal 155 |
| crystallization-induced emission | energy level 124, 171, 198 |
| enhancement 191 | continuous 43, 125 |
| | discrete 43, 124, 125, 156, 175, |
| density functional perturbation | 207 |
| theory 294 | dispersed 166 |
| density functional theory (DFT) | ground-state 198 |
| 99, 127, 170, 171, 197, 199, | EPR see electron paramagnetic |
| 232, 238, 268, 285, 286, 288, | resonance |
| 290-292, 294, 296, 316-318 | ESA see excited state absorption |
| DFT see density functional theory | excited state absorption (ESA) |
| dopant 238, 275, 298 | 157, 158, 178, 199 |
| | |

Faradaic efficiency 272 grand unified model (GUM) fluorescence 173, 184, 193, 129-136, 138 195-197 ground-state bleaching (GSB) 156, Fourier transform 275 157, 178 Fourier transform infrared GSB see ground-state bleaching spectroscopy (FTIR) 237-239, GSB band 156 245, 275 GSB peak 158 FTIR see Fourier transform GSB signal 157, 158, 200 infrared spectroscopy GUM see grand unified model gap 43, 62, 98, 124, 135, 150, 161, highest occupied molecular orbital 166, 234, 250, 293 (HOMO) 4, 125, 150, 152-154, gauge-independent atomic orbital 166, 171-174, 177, 184, 293, 294 295, 303 gel electrophoresis 5, 17, 18, 232 high-speed mechanical shutter gold cluster 173, 175, 176, 198 180-182, 185, 197, 199, HOMO see highest occupied 201–205, 207, 208, 229, 230, molecular orbital hydrogenation 230, 232, 233, 243, 232, 234, 236–238, 247, 249 chiral origin of 203 265-268, 270, 274, 312, 314, homochiral 206 315 ligand-protected 131, 175, 205 stereoselective 262, 263 ligand-stabilized 175 hydroxyapatite 262, 266 luminescence of 181, 184, 191 hyperpolarizability 195, 197 optical stability of 202, 208 ultrasmall 196 imaging 195 gold nanocluster (AuNC) 5-7, biological 194, 196 high-resolution multiphoton 13-15, 25-27, 43, 44, 61, 62, 80, 81, 110, 123-138, 166, 166, 193, 207 live cell 195 168–170, 176–182, 186–188, 192–194, 196, 198–206, 231, low-power medical 193 second harmonic 196 232, 260, 261, 265 isolation 4, 5, 14, 15, 17, 20, 30, gold nanoparticle (AuNP) 98, 99, 101, 103, 105-111, 113, 114, 71, 178 149, 150, 166–170, 176, 177, chromatographic 60 180, 196, 199, 230, 231, 236, gel 6 237, 240 subtle 4, 5, 29 gold nanorod 168, 233 isomer 25, 27, 99-103, 107, 109, gold nanowire 168 110, 206, 270 GPC see gel permeation dextrorotatory 204 chromatography levorotatory 204

optical 205 structural 12, 13, 191 isomerization 296, 304-306

Jellium model 128, 129

kernel 70, 75, 104, 105, 109, 113, 131, 132, 135-138, 201, 204, 271, 279 bi-icosahedron 108 double helix 113 high-symmetry 108 limited-sized 136 metal 70 reasonable high-symmetry 111 tetrahedron 113 Kubo criterion for metallic state 157, 161

Langmuir-Hinshelwood mechanism 125 laser 178, 192, 196, 198 infrared 195 near-infrared 196 pulsed 198 laser desorption/ionization mass spectrometry (LDIMS) 14, 20 laser fluence 158, 178, 179 laser mode-locking technology 198 LDIMS see laser desorption/ ionization mass spectrometry LEIST see ligand-exchange-induced size/structure transformation Lewis acid 237, 243, 244, 264, 265 ligand 24, 69-73, 78-82, 87, 88, 110, 113, 114, 129, 130, 180-182, 189, 197, 201, 205, 257, 258, 260, 265-268 ligand exchange 14, 24, 48, 72, 73,

98, 182, 197, 296, 311

ligand-exchange-induced size/ structure transformation (LEIST) 14, 47, 48, 53, 60 localized surface plasmon resonance 176 lowest unoccupied molecular orbital (LUMO) 4, 150, 152-154, 166, 171-173, 177, 184, 293, 295, 301, 303 luminescence 184, 185, 188-191, 201, 207 aggregation-induced 207 near-infrared 181, 182 LUMO see lowest unoccupied molecular orbital

Mackay icosahedron (MI) 49,50 MALDIMS see matrix-assisted laser desorption ionization mass spectrometry matrix-assisted laser desorption ionization mass spectrometry (MALDIMS) 8, 20, 25, 46, 231 Maxwell equation 169 MI see Mackay icosahedron Mie theory 169 model 42, 127-130, 138, 279, 287, 292, 297, 314 anion adsorption 299 charge distribution 301 continuum solvation 311 motif 49, 77-79, 81, 86, 108, 110, 127, 128

nanocatalysis 230, 250, 257 nanocatalyst 230, 245, 255, 256 nanocluster 41-44, 46-49, 52-54, 59-62, 72-87, 97, 98, 108–110, 114, 155, 156, 178, 179, 191–194, 232, 233, 255, 256, 258, 259, 261, 262, 264-268, 270, 271, 275, 277, 280

| atomically precise 45, 47, 151 | simulated isotope 19 |
|--------------------------------------|------------------------------------|
| bimetal 261 | stripe 57 |
| excitonic 63, 155, 156 | surface helical 49 |
| fcc type 131 | surface protection 63 |
| polydisperse 45, 55 | PES see potential energy surface |
| thiolate-protected 264 | phenomenon 7, 76, 82, 105, 167, |
| nanoparticle 41, 42, 44-46, 48, | 168, 174, 188, 189, 193, 197, |
| 50, 52, 98, 150, 151, 155, 156, | 202, 262 |
| 167-169, 175, 176, 178, 179, | life 202 |
| 193-195, 202, 256, 267, 268 | luminescence enhancement 191 |
| atomically precise 154 | nonlinear optical 196 |
| metallic-state 155 | rainbow 167 |
| plasma 178 | second harmonic 196 |
| ultrasmall 149 | phonon 155, 156, 160, 200 |
| NMR see nuclear magnetic | photocatalysis 240, 241, 273, 295 |
| resonance | photoexcitation 155, 157, 158, |
| nuclear magnetic resonance | 160, 178, 187 |
| (NMR) 8, 182, 184, 240, 242, | photoluminescence 4, 70, 166, |
| 294 | 180, 185, 207 |
| | photon 193, 195 |
| optical activity 166, 197, 202, 205, | photo-oxidation 233, 241, 242 |
| 206 | photosensitizer 233, 234, 236 |
| optical properties 149–155, 157, | plasmon 151, 162, 199, 200 |
| 159, 165–169, 171, 173–175, | polarization coefficient 192 |
| 177–185, 187, 189, 191, 196, | polyacrylamide gel electrophoresis |
| 203, 205, 207, 285 | (PAGE) 5, 17–20, 27, 232 |
| optical spectrum 108, 110, 153 | potential energy surface (PES) |
| photocatalytic 236 | 99–102, 106, 110, 269, 271, |
| photoculary tro | 276, 291, 295, 296, 302, 312, |
| PAGE see polyacrylamide gel | 314 |
| electrophoresis | prediction 24, 41, 100, 106, 111, |
| pattern 104, 105, 108, 110 | 114, 128, 291 |
| carved 166 | quantitative 317 |
| cyclic tetramer 54 | preparative thin-layer |
| divide and protect 107 | chromatography (PTLC) 27, |
| face-centered cubic stacking | 29 |
| 113 | |
| growth 103, 106, 112-114 | quantum confinement effect 4, |
| herringbone 54 | 165, 166, 169, 207, 256, 272 |
| interlocked staple motif 110 | quantum size effect 125, 150 |
| monomeric staple motif 51 | • |
| novel thiol bonding 126 | radial distribution function 292 |

| cluster-catalyzed 296 heterogeneous 229–231, 260, 265 hydrogen evolution 274, 300 photocatalytic 241 reduction 6–8, 46, 187, 207, 273, 308 anti-galvanic 28 controlled 59 electrocatalytic 271 | spectrum 9, 19, 25, 28, 151, 152, 154, 155, 157, 158, 171, 177, 179, 200, 293, 297 diffuse reflectance UV-Vis 240 theoretical circular dichroism 108 SPR see surface plasmon resonance staple 25, 43, 50, 271, 274, 286, 287, 307, 308 dethiolated 311 |
|--|---|
| resonance 176 electron paramagnetic 258, 294 nuclear magnetic 182, 240, 294 nuclear paramagnetic 8 | ionic 271, 308 pentameric 74, 75 tetrameric 153 trimeric 54 superatom complex model (SAC |
| SAM <i>see</i> self-assembled monolayer scanning tunneling microscopy 126 | model) 129, 138 superatom network model (SAN model) 129, 138 |
| second harmonic generation (SHG) 196, 197 selective aerobic oxidation 230, 233, 234 | surface plasmon resonance (SPR) 70, 150, 151, 154, 155, 166, 168–169, 175, 177, 199 surface protection 43, 44, 51–53, |
| self-assembled monolayer (SAM) 41, 126 | 55, 56, 58, 60–62 |
| semihydrogenation 245–247, 265, 266 | TDDFT see time-dependent density functional theory |
| separation 4, 5, 14, 15, 17, 20, 22–24, 26–29, 203, 205, 208, 273, 286 electron–hole 273 enantiomeric 205 high-resolution 21, 22 | capillary electrophoretic 29 high-resolution isolation 5 hyper-Rayleigh scattering 195 liquid chromatography 21 mass spectrometric 6 molecular 42 |
| SHG see second harmonic generation | pump-probe 198 spectroscopic 294 |
| Slater–Janak transition state method 294 active oxygen 258, 259, 261, 273 electron-deficient oxygen 262 electronegative 265 optically active 205, 256 potential isoelectronic 130 | thiol 5, 7, 10, 12, 46, 47, 55–57, 126, 175, 176, 178, 180, 193, 196, 199, 303, 304 thiolate 5, 13, 69–71, 73, 74, 77, 104, 106, 108, 188, 189, 258, 260, 261, 264, 266, 267, 306–308, 310 achiral 26 |

bridging 58, 62, 204 dangling 311 residual 259 terminal 107, 109 thiolate complex 180, 181, 187-189 time-dependent density functional theory (TDDFT) 170, 171, 174, 294, 295 TPA see two-photon absorption TPF see two-photon fluorescence transition 62, 150, 152, 154, 155, 161, 171, 173-179, 193, 196, 198-200, 207, 304, 305 excitonic 157 in-band 171 interband 152, 173, 174 metal-to-molecule 199

metal-to-nonmetal 162 single-electron 154 transition state (TS) 101, 293-296, 304-306 transition state theory 295, 296 TS see transition state two-photon absorption (TPA) 192-197, 207 two-photon fluorescence (TPF) 193, 194, 197, 207

ultraviolet photoemission spectroscopy (UPS) 236, 294 UPS see ultraviolet photoemission spectroscopy

X-ray crystallography 5, 9, 42, 52, 62, 76, 80

"As no two identical leaves exist in this world, similarly, no two identical nanoparticles exist in a solution. Fortunately, atomically precise nanoclusters with absolute purity have alleviated this problem to some extent. This well-written book systematically overviews the synthesis, structural determination, optical properties, and catalysis applications of nearly 150 such known nanoclusters. The personal experiences of the authors and their pioneer perspectives inspire to explore the deep sea of unknowns of these nanoclusters."

Prof. Zhiyong Tang National Center for Nanoscience and Technology, China

"One of the foremost challenges of nanoscience is to develop atomic-size structures and then establish precise structure—property correlations at the atomic and even at the single-electron level. This splendid book thoroughly reviews the latest aspects of metal nanoclusters and is highly recommended as a desktop reference for senior scientists and an enlightening introductory book for young scientists, students, and those new to nanoclusters."

Prof. Di Sun Shandong University, China

"This is a timely book on the fascinating world of metal nanoclusters and covers the methods of synthesis and structural determination of ligand-protected metal nanoclusters and their optical and catalytic properties. It is greatly beneficial to communities of cluster chemistry, nanoscience, and nanotechnology."

Prof. Quan-Ming Wang Tsinghua University, China

The primary goal of nanotechnology is to achieve nanoscale materials and devices with atomic precision. This book presents the recent breakthroughs in the solution-phase synthesis and applications of atomically precise nanoclusters and the exciting progress in this new research field. The chapters are contributed by leading experts in the field and cover the methods of synthesis, atomic structures, electronic and optical properties, and catalytic applications of noble metal nanoclusters. The book highlights the exciting opportunities these new nanocluster materials offer to chemists and physicists to enhance their understanding of the fundamental science of nanoclusters, including atomic-level structure–property correlation and design of new materials, as well as their applications, including catalysis, biomedicine, sensing, imaging, optics, and energy conversion. The book is of interest to readers and researchers in nanotechnology, nanochemistry, catalysis, and computational chemistry, as well as practitioners in industry R&D for new materials. It is written to be accessible to undergraduate and graduate students and, therefore, is an excellent teaching material.



Yan Zhu is a full professor of physical chemistry at Nanjing University, China. Her research interests focus on the design of atomically precise metal nanoclusters as model catalysts for catalytic applications, including thermocatalysis, photocatalysis, and electrocatalysis science.



Rongchao Jin is a professor of chemistry at Carnegie Mellon University, USA. He earned his PhD in chemistry from Northwestern University, USA, in 2003. He joined the chemistry faculty of Carnegie Mellon University in 2006 and was promoted to associate professor in 2012 and full professor in 2015. His research interests include atomically precise nanoparticles, optics of nanoparticles, and catalysis.



