

## Bibliography

- Amer, N. M., Skumanich, A. and Jackson, W. B. (1983). The contribution of the Staebler–Wronski effect to gap-state absorption in hydrogenated amorphous silicon, *Physica*, **117 B**, pp. 897–898.
- Aoki, T., Ikeda, K., Kobayashi, S. and Shimakawa, K. (2006). Recombination kinetics of very-long-lived photoluminescence decay in a-Si:H correlated with light-induced spin densities, *Philos. Mag. Lett.*, **86**, pp. 137–145.
- Aoki, T., Komodoori, S., Kobayashi, S., Fujihashi, C., Ganjoo, A. and Shimakawa, K. (2002). Photoluminescence lifetime distribution of a-Si:H and a-Ge:H expanded to nanosecond region using wide-band frequency-resolved spectroscopy, *J. Non-Cryst. Solids*, **299–302**, pp. 642–647.
- Astakov, O., Carius, R., Lambertz, A., Petrusenko, Yu., Borysenko, V., Barankov, D. and Finger, F. (2008). Structure of the ESR spectra of thin film silicon after electron bombardment, *J. Non-Cryst. Solids*, **354**, pp. 2329–2332.
- Biegelsen, D. K. and Street, R. A. (1980). Photoinduced defects in chalcogenide glasses, *Phys. Rev. Lett.*, **44**, pp. 803–806.
- Bishop, S. G., Strom, U. and Taylor, P. C. (1977). Optically induced metastable paramagnetic states in amorphous semiconductors, *Phys. Rev. B*, **15**, pp. 2278–2294.
- Biswas, R. and Kwon, I. (1991) *New interpretations of the Staebler-Wronski effect in a-Si:H with molecular dynamics simulations*, AIP Conf. Proc. (American Inst. Phys. New York), pp. 45–50.
- Boehme, C. and Lips, K. (2003). Theory of time-domain measurement of spin-dependent recombination with pulsed electrically detected magnetic resonance, *Phys. Rev. B*, **68**, pp. 245105-1–245105-19.
- Boulitrop, F., Chenevas-Paule, A. and Dunstan, D. J. (1983). Luminescence and magnetic resonance in post-hydrogenated microcrystalline silicon, *Solid State Commun.*, **48**, pp. 181–184.
- Branz, H. (1999). Hydrogen collision model: Quantitative description of metastability in amorphous silicon, *Phys. Rev. B*, **59**, pp. 5498–5512.
- Branz, H. (2003). The hydrogen collision model of metastability after 5 years: experimental tests and theoretical extensions, *Sol. Energy Mater. & Sol. Cells*, **78**, pp. 425–445.

- Brower, K. L. (1983).  $^{29}\text{Si}$  hyperfine structure of unpaired spins at the Si/SiO<sub>2</sub> interface, *Appl. Phys. Lett.*, **43**, pp. 1111–1113.
- Brower, K. L. (1986). Strain broadening of the dangling-bond resonance at the (111) Si-SiO<sub>2</sub> interface, *Phys. Rev. B*, **33**, pp. 4471–4478.
- Butté, R., Vignoli, S., Meaudre, M., Meaudre, R., Marty, O., Savior, L. and Roca i Cabarrocas, P. (2000). Structural, optical and electronic properties of hydrogenated polymorphous silicon films deposited at 150°C, *J. Non-Cryst. Solids*, **266–269**, pp. 263–268.
- Caputo, D., de Cesare, G., Irrera, F, Palma, F, Rossi, M. C., Conte, G., Nobile, G. and Fameli, G. (1994). A systematic investigation of the role of material parameters in metastability of hydrogenated amorphous silicon, *J. Non-Cryst. Solids*, **170**, pp. 278–286.
- Carlson, D. E. (1986). Hydrogenated microvoids and light-induced degradation of amorphous-silicon solar cells, *Appl. Phys. A*, **41**, pp. 305–309.
- Cavenett, B. C. (1981). Optically detected magnetic resonance (O. D. M. R) investigations of recombination processes in semiconductors, *Adv. Phys.*, **30**, pp. 475–538.
- Cavenett, B. C., Depinna, S. P., Austin, I. G., and Searle, T. M. (1983). Determination of the g values of the ODMR signals in a-Si:H, *Philos. Mag. B*, **48**, pp. 169–185.
- Chadi, D. J. and Chang, K. J. (1989). Energetics of DX-center formation in GaAs and Al<sub>x</sub>Ga<sub>1-x</sub>As alloys, *Phys. Rev. B*, **39**, pp. 10063–10074.
- Cody, G. D., Tiedje, T., Abeles, B., Brooks, B. and Goldstein, Y. (1981). Disorder and the optical-absorption edge of hydrogenated amorphous silicon, *Phys. Rev. Lett.*, **47**, pp. 1480–1483.
- Cohen, M. H., Fritzsche, H. and Ovshinsky, S. R. (1969). Simple band model for amorphous semiconductors alloys, *Phys. Rev. Lett.*, **22**, pp. 1065–1068.
- Collins, R. W. and Paul, W. (1982). Model for the temperature dependence of photoluminescence in a-Si:H and related materials, *Phys. Rev. B*, **25**, pp. 5257–5262.
- Das, D., Shirai, H., Hanna, J. and Shimizu, I. (1991). Narrow band-gap a-Si:H with improved minority carrier-transport prepared by chemical annealing, *Jpn. J. Appl. Phys.*, **30**, pp. 239–242.
- de Lima, Taylor, P. C., Morrison, S, LeGeune, A. and Marques, F. C. (2002). ESR observations of paramagnetic centers in intrinsic hydrogenated microcrystalline silicon, *Phys. Rev. B*, **65**, pp. 235324-1–235324-6.

- Depinna, S. P., Homewood, K., Cavenett, B. C., Austin, I. G., Searle, T. M., Willeke, G. and Kinmond, S. (1983). O.D.M.R. investigation of recombination in  $\mu\text{c-Si:H}$ , *Philos. Mag. B*, **47**, pp. L57–L62.
- Depinna, S. P. and Dunstan, D. J. (1984). Frequency-resolved spectroscopy and its application to the analysis of recombination in semiconductors, *Philos. Mag. B*, **50**, pp. 579–597.
- Dersch, H., Stuke, J. and Beichler, J. (1981). Light-induced dangling bonds in hydrogenated amorphous silicon, *Appl. Phys. Lett.*, **38**, pp. 456–458.
- Dunstan, D. J. and Boulitrop, F. (1984). Photoluminescence in hydrogenated amorphous silicon, *Phys. Rev. B*, **30**, pp. 5945–5957.
- Ehara, T., Ikoma, T. and Tero-Kubota, S. (2000). Multi-band electron paramagnetic resonance study of the defects in microcrystalline silicon, *J. Non-Cryst. Solids*, **266–269**, pp. 540–543.
- Elliott, S. R. and Shimakawa, K. (1990). Model for bond-breaking mechanisms in amorphous arsenic chalcogenides leading to light-induced electron-spin resonance, *Phys. Rev. B*, **42**, pp. 9766–9770.
- Engemann, D. and Fischer, R. (1973). Radiative recombination in amorphous silicon, *Proc. 5th Int. Conf. on Amorphous and Liquid Semiconductors*, pp. 947–952.
- Engemann, D. and Fischer, R. (1974). Influence of preparation conditions on the radiative recombination in amorphous silicon, *Proc. 12th Int. Conf. on the Physics of Semiconductors*, pp. 1042–1046.
- Englman, R. and Jortner, J. (1970). The energy gap law for radiative transitions in large molecules, *Mol. Phys.*, **18**, pp. 145–164.
- Fehr, M., Schnegg, A., Rech, B., Lips, K., Astakov, O., Finger, F., Pfanner, G., Freysoldt, C., and Neugebauer, J., Bittl, R., and Teutloff, C. (2011). Combined multifrequency EPR and DFT study of dangling bonds in a-Si:H, *Phys. Rev. B*, **84**, pp. 245203-1–10.
- Fehr, M., Schnegg, A., Teutloff, C., Bittl, R., Astakhov, O., Finger, F., Rech, B. and Lips, K. (2009). Hydrogen distribution in the vicinity of dangling bonds in hydrogenated amorphous silicon (a-Si:H), *Phys. Status Solidi A*, **207**, pp. 552–555.
- Finger, F. (2004). Defects and structure of hydrogenated microcrystalline silicon films deposited by different techniques, *J. Non-Cryst. Solids*, **338–340**, pp. 168–172.
- Finger, F., Carius, R., Dylla, T., Klein, S., Okur, S. and Günes, M. (2005). Instability phenomena in microcrystalline silicon films, *J. Optoelectron. Adv. Mat.*, **7**, pp. 83–90.

- Finger, F., Malten, C., Hapke, P., Carius, R., Flückiger, R. and Wagner, H. (1994). Free electrons and defects in microcrystalline silicon studied by electron spin resonance, *Philos. Mag. Lett.*, **70**, pp. 247–254.
- Finger, F., Müller, J., Malten, C. and Wagner, H. (1998). Electronic states in hydrogenated microcrystalline silicon, *Philos. Mag. B*, **77**, pp. 805–830.
- Fontenberta, i Morral, A., Brenot, R., Hamers, E. A. G., Vanderhaghen, R. and Roca i Cabarrocas, P. (2000). In situ investigation of polymorphous silicon deposition, *J. Non-Cryst. Solids*, **266–269**, pp. 48–53.
- Freysoldt, C., Pfanner, G. and Neugebauer, J. (2012). The dangling-bond defect in amorphous silicon: statistical random versus kinetically derived defect geometries, *J. Non-Cryst. Solids*, **358**, pp. 2063–2066.
- Fritzsche, H. (2001). Development in understanding and controlling the Staebler-Wronski effect in a-Si:H, *Annu. Rev. Mater. Res.*, **31**, pp. 47–79.
- Fuhs, W., Mell, H., Stuke, J., Thomas, P. and Weiser, G. (1985). Photo-induced metastable effects in hydrogenated amorphous silicon (a-Si:H), *Ann. Phys.*, **42**, pp. 187–197.
- Fujita, Y., Yamaguchi, M. and Morigaki, K. (1994). Kinetics of annealing of dangling bonds in sputtered amorphous silicon and hydrogenated amorphous silicon, *Philos. Mag. B*, **69**, pp. 57–67.
- Ganjoo, A., Ikeda, Y. and Shimakawa, K. (2000). In situ measurements of photo-induced volume changes in amorphous chalcogenide films, *J. Non-Cryst. Solids*, **266–269**, pp. 919–923.
- Gleskova, H., Morin, P. A. and Wagner, S. (1993). Kinetics of recovery of the light-induced defects in hydrogenated amorphous silicon under illumination, *Appl. Phys. Lett.*, **62**, pp. 2063–2065.
- Godet, C. (1998). Metastable hydrogen atom trapping in hydrogenated amorphous silicon films: a microscopic model for metastable defect creation, *Philos. Mag. B*, **77**, pp. 765–777.
- Godet, C., Morin, P. and Roca i Cabarrocas, P. (1996). Influence of the dilute-phase SiH bond concentration on the steady-state defect density in a-Si:H, *J. Non-cryst. Solids*, **198–200**, pp. 449–452.
- Graeff, C. F. O., Buhleier, R. and Stutzmann, M. (1993). Light-induced annealing of metastable defects in hydrogenated amorphous silicon, *Appl. Phys. Lett.*, **62**, pp. 3001–3003.
- Guéron, M. and Solomon, I. (1965). Effect of spin resonance on hot electrons by spin-orbit coupling in n-type InSb, *Phys. Rev. Lett.*, **15**, pp. 667–670.
- Han, D., Yoshida, M. and Morigaki, K. (1987). Light-induced metastable defects in a-Si:H elucidated by optically detected magnetic resonance measurements at 2 K, *Solid State Commun.*, **63**, pp. 1083–1086.

- Hautala, J., Ohlsen, W. D. and Taylor, P. C. (1988). Optically induced electron-spin resonance in  $\text{As}_x\text{S}_{1-x}$ , *Phys. Rev. B*, **38**, 11048–11060.
- Hikita, H., Takeda, K., Kimura, Y., Yokomichi, H. and Morigaki, K. (1997). Deconvolution of ESR spectra and their light-induced effect in a-Si:H, *J. Phys. Soc. Jpn.*, **66**, pp. 1730–1740.
- Hikita, H., Takeda, K., Yokomichi, H. and Morigaki, K. (unpublished).
- Hirabayashi, I., Morigaki, K. and Nitta, S. (1980). New evidence for defect creation by high optical excitation in glow discharge amorphous silicon, *Jpn. J. Appl. Phys.*, **19**, pp. L357–L360.
- Hirose, T., Yoshioka, H. and Horai, K. (1966). Conductivity increase of reduced rutile with the saturation of electron spin resonance, *J. Phys. Soc. Jpn.*, **21**, pp. 559–559.
- Honig, A. (1966). Neutral-impurity scattering and impurity Zeeman spectroscopy in semiconductors using highly spin-polarized carriers, *Phys. Rev. Lett.*, **17**, 186–188.
- Ishii, N., Kumeda, M. and Shimizu, T. (1981). The  $g$ -value of defects in amorphous C, Si and Ge. *Jpn. J. Appl. Phys.*, **20**, L673–L676.
- Ishii, N. and Shimizu, T. (1998). Cluster-model calculations of hyperfine parameters and  $g$ -values for defects in a-Si:H, *J. Non-Cryst. Solids*, **227–230**, pp. 358–361.
- Isoya, J., Yamasaki, S., Okushi, H., Matsuda, A. and Tanaka, K. (1993). Electron-spin-echo envelope-modulation study of the distance between dangling bonds and hydrogen atoms in hydrogenated amorphous silicon, *Phys. Rev. B*, **47**, pp. 7013–7024.
- Jackson, W. B. and Kakalios, J. (1988). Evidence for hydrogen motion in annealing of light-induced metastable defects in hydrogenated amorphous silicon, *Phys. Rev. B*, **37**, pp. 1020–1023.
- Jones, R. and Lister, G. M. S. (1990). *Ab initio* calculations on metastable defects in a-Si:H. The Staebler-Wronski effect., *Philos. Mag. B*, **61**, pp. 881–894.
- Kakalios, J., Street, R. A. and Jackson, W. B. (1987). Stretched-exponential relaxation arising from dispersive diffusion of hydrogen in amorphous silicon, *Phys. Rev. Lett.*, **59**, pp. 1037–1040.
- Kamei, T., Hata, N., Matsuda, A., Uchiyama, T., Amano, S., Tsukamoto, K., Yoshioka, Y. and Hirao, T. (1996). Deposition and extensive light soaking of highly pure hydrogenated amorphous silicon, *Appl. Phys. Lett.*, **68**, 2380–2382.
- Kamimura, H. and Mott, N. F. (1976). The variable range hopping induced by electron spin resonance in n-type silicon and germanium, *J. Phys. Soc. Jpn.*, **40**, pp. 1351–1358.

- Kaplan, D., Solomon, I. and Mott, N. F. (1978). Explanation of the large spin-dependent recombination effect in semiconductors, *J. de Physique*, **39**, pp. L51–L54.
- Kastner, M., Adler, D. and Fritzsche, H. (1976). Valence-alternation model for localized gap states in lone-pair semiconductors, *Phys. Rev. Lett.*, **37**, pp. 1504–1507.
- Kimerling, L. C. (1978). Recombination enhanced defect reactions, *Solid State Electron.*, **21**, pp. 1391–1401.
- Kishimoto, N. and Morigaki, K. (1977). Resistivity decrease due to electron spin resonance in the metallic region of heavily phosphorus-doped silicon, *J. Phys. Soc. Jpn.*, **42**, pp. 137–145.
- Kishimoto, N., Morigaki, K. and Murakami, K. (1981). Conductivity change due to electron spin resonance in amorphous Si-Au system, *J. Phys. Soc. Jpn.*, **50**, pp. 1970–1977.
- Kondo, M. and Morigaki, K. (1990). Light-induced phenomena in a-Si:H as elucidated by optically detected electron nuclear double resonance, *Proc. Int. Conf. Physics of Semiconductors*, eds. Anastassakis, E. M. and Joannopoulos, J. D. (World Scientific, Singapore), pp. 2083–2086.
- Kondo, M. and Morigaki, K. (1991). The role of hydrogen clusters in the Staebler-Wronski effect of amorphous silicon as elucidated by optically detected electron nuclear double resonance, *J. Non-Cryst. Solids*, **137–138**, pp. 247–250.
- Kondo, M. and Morigaki, K. (1993). Possibility of hydrogen migration in photoinduced defect creation process of a-Si:H, *J. Non-Cryst. Solids*, **164–166**, pp. 227–230.
- Kondo, M., Nishimiya, T., Saito, K. and Matsuda, A. (1998). Light induced phenomena in microcrystalline silicon, *J. Non-Cryst. Solids*, **227–230**, pp. 1031–1035.
- Kondo, M., Yamasaki, S. and Matsuda, A. (2000). Microscopic structure of defects in microcrystalline silicon, *J. Non-Cryst. Solids*, **266–269**, pp. 544–547.
- Kugler, S. (2012). Advances in understanding the defects contributing to the tail states in pure amorphous silicon, *J. Non-Cryst. Solids*, **358**, pp. 2060–2062.
- Lang, D. V. (1992). *Deep Centers in Semiconductors: A State-of-Art Approach*, S. T. Chapter 7, “DX Centers in III–V Alloys” (Gordon and Breach Science Publishers, Switzerland), pp. 591–642.
- Lang, D. V. and Kimerling, L. C. (1974). Observation of recombination-enhanced defect reactions in semiconductors, *Phys. Rev. Lett.*, **33**, pp. 489–492.

- Lang, D. V. and Kimerling, L. C. (1976). Observation of athermal defect annealing in GaP, *Appl. Phys. Lett.*, **28**, pp. 248–250.
- Lang, D. V. and Logan, R. A. (1977). Large-lattice-relaxation model for persistent photoconductivity in compound semiconductors, *Phys. Rev. Lett.*, **39**, pp. 635–639.
- Lang, D. V. and Logan, R. A. (1979). *Physics of Semiconductors*, 1978, Conference Series No. 43, ed. Wilson, B. L. H., Chapter 13 “Defects Chemical Shifts of DX Centers in  $\text{Al}_x\text{Ga}_{1-x}\text{As}$ ” (Institute of Physics, Bristol and London), pp. 433–436.
- Lepine, D. (1972). Spin-dependent recombination on silicon surface, *Phys. Rev. B*, **6**, pp. 436–441.
- Lepine, D., Grazhulis, V. A., and Kaplan, D. (1976). Spin-dependent recombination on dislocation in silicon, *Proc. 13th Int. Conf. Phys. Semicond.*, Rome, ed. Fumi, E. G., pp. 1081–1084.
- Lips, K., Boehme, C. and Ehara, T. (2005). The impact of the electron spin on charge carrier recombination: the example of amorphous silicon, *J. Optoelec. Adv. Mat.*, **7**, pp. 13–24.
- Lips, K., Kanschä, P. and Fuhs, W. (2003). Defects and recombination in microcrystalline silicon, *Sol. Energy Mat. & Sol. Cells*, **78**, pp. 513–541.
- Longeaud, C., Roy, D. and Saadane, O. (2002). Role of interstitial hydrogen and voids in light-induced metastable defect formation in hydrogenated amorphous silicon: A model, *Phys. Rev. B*, **65**, pp. 085206-1–085206-9.
- Lukács, R. and Kugler, S. (2011). Photoinduced volume changes of obliquely and flatly deposited amorphous AsSe films universal description of the kinetics, *Jpn. J. Appl. Phys.*, **50**, pp. 091401-1–091401-4.
- Lukács, R., Veres, M., Shimakawa, K. and Kugler, S. (2010). On photoinduced volume change in amorphous selenium: Quantum chemical calculation and Raman spectroscopy, *J. Appl. Phys.*, **107**, pp. 073517-1–073517-5.
- Malten, C., Carius, R., Finger, F. and Yamasaki, S. (1998). ODMR measurements of microcrystalline silicon, *MRS Proc.*, **507**, pp. 787–792.
- Maxwell, R. and Honig, A. (1966). Neutral-impurity scattering experiments in silicon with highly spin-polarized electrons, *Phys. Rev. Lett.*, **17**, pp. 188–190.
- Matsuda, A. (2006) *Springer Handbook of Electronic and Photonic Materials*, eds. Kasap, S. and Capper, P., Part C 26 “amorphous and Microcrystalline Silicon” (Springer, New York), pp. 581–595.
- Mooney, P. M. (1992). *Deep Centers in Semiconductors: A State-of-Art Approach*, ed. Pantelides, S. T., Chapter 7, “DX Centers in III–V Alloys:

- Recent Developments" (Gordon and Breach Science Publishers, Switzerland), pp. 643–665.
- Morigaki, K. (1981). Spin-dependent radiative and nonradiative recombinations in hydrogenated amorphous silicon: Optically detected magnetic resonance, *J. Phys. Soc. Jpn.*, **50**, pp. 2279–2287.
- Morigaki, K. (1983). Optically detected magnetic resonance in amorphous semiconductors, *Jpn. J. Appl. Phys.*, **22**, pp. 375–388.
- Morigaki, K. (1984) *Semiconductors and Semimetals 21 Part C*, ed. Pankove, J. I., Chapter 4 "Optically Detected Magnetic Resonance" (Academic Press, Orlando), pp. 155–191.
- Morigaki, K. (1988). Microscopic mechanism for the photo-creation of dangling bonds in a-Si:H, *Jpn. J. Appl. Phys.*, **27**, pp. 163–168.
- Morigaki, K. (1992). Self-trapping of holes and light- and thermally induced defect creation in a-Si:H, *J. Non-Cryst. Solids*, **141**, pp. 166–175.
- Morigaki, K. (1999) *Physics of Amorphous Semiconductors* (World Scientific, Singapore, and Imperial College Press, London).
- Morigaki, K. (2001). Role of hydrogen-related weak bonds and hydrogen-related dangling bonds in light-induced structural changes in a-Si:H, *Bull. Hiroshima Inst. Tech.*, **35**, pp. 47–56.
- Morigaki, K. (2002a). Photoluminescence in amorphous solids, *Nonlinear Opt.*, **29**, pp. 265–272.
- Morigaki, K. (2002b). On the electron spin resonance of band tails and defects in hydrogenated amorphous silicon, *Bull. Hiroshima Inst. Tech.*, **36**, pp. 25–30.
- Morigaki, K. (2009). Recombination processes and light-induced defect creation in hydrogenated amorphous silicon, *Phys. Status Solidi A*, **206**, pp. 868–873.
- Morigaki, K. and Hikita, H. (1999). Hydrogen-related dangling bonds in hydrogenated amorphous silicon, *Bull. Hiroshima Inst. Tech.*, **33**, pp. 135–142.
- Morigaki, K. and Hikita, H. (2005). Light-induced creation of hydrogen-pairs in a-Si:H, *Solid State Commun.*, **136**, pp. 616–620.
- Morigaki, K. and Hikita, H. (2007). Modeling of light-induced defect creation in hydrogenated amorphous silicon, *Phys. Rev. B*, **76**, pp. 085201-17.
- Morigaki, K. and Hikita, H. (2011). Stretched exponential relaxation processes in hydrogenated amorphous and polymorphous silicon, *Phys. Status Solid C*, **8**, pp. 2564–2568.
- Morigaki, K. and Hikita, H. (unpublished).



- Morigaki, K., Hikita, H. and Kondo, M. (1995). Self-trapping of holes and related phenomena in a-Si:H, *J. Non-Cryst. Solids*, **190**, pp. 38–47.
- Morigaki, K., Hikita, H. and Ogihara, C. (2009a). Light-induced defect creation under intense pulsed illumination in hydrogenated amorphous silicon, *J. Optoelec. Adv. Mater.*, **11**, pp. 1–14.
- Morigaki, K., Hikita, H., Takeda, K. and Roca I Cabarrocas, P. (2010). The kinetics of the light-induced defect creation in hydrogenated polymorphous silicon—stretched exponential relaxation, *Phys. Status Solidi C*, **7**, pp. 692–695.
- Morigaki, K., Hikita, H., Takemura, H., Yoshimura, T. and Ogihara, C. (2003). Light-induced defect creation under pulsed subbandgap illumination in hydrogenated amorphous silicon, *Philos. Mag. Lett.*, **83**, pp. 341–349.
- Morigaki, K., Hikita, H., Yamaguchi, M. and Fujita, Y. (1998). The structure of dangling bonds having hydrogen at a nearby site in a-Si:H, *J. Non-Cryst. Solids*, **227–230**, pp. 338–342.
- Morigaki, K., Hikita, H., Yamaguchi, M. and Fujita, Y. (2003). Anisotropic magnetic centers and conduction electrons in hydrogenated microcrystalline silicon, *Mat. Eng., B*, **103**, pp. 37–44.
- Morigaki, K., Hirabayashi, I., Nakayama, M., Nitta, S. and Shimakawa, K. (1980). Fatigue effect in luminescence of glow discharge amorphous silicon at low temperatures, *Solid State Commun.*, **33**, pp. 851–856.
- Morigaki, K., Kishimoto, N. and Lepine, D. J. (1975). Resistivity decreases due to electron spin resonance in the metallic region of heavily phosphorus doped silicon, *Solid State Commun.*, **17**, pp. 1017–1019.
- Morigaki, K. and Kondo, M. (1995). Optically detected magnetic resonance in a-Si:H, *Solid State Phenomena*, **44–46**, pp. 731–764.
- Morigaki, K. and Niikura, C. (2005). The *g*-values of defects in hydrogenated microcrystalline silicon, *Solid State Commun.*, **136**, pp. 308–312.
- Morigaki, K., Niikura, C., Bourée, J. E. and Equer, B. (2002). Anisotropic magnetic centres in hydrogenated microcrystalline and polymorphous silicon, *J. Non-Cryst. Solids*, **299–302**, pp. 561–565.
- Morigaki, K., Niikura, C., Hikita, H. and Yamaguchi, M. (2009b). Defects in hydrogenated microcrystalline silicon prepared by plasma-enhanced chemical vapour deposition, *J. Appl. Phys.*, **105**, 083703-1–083703-6.
- Morigaki, K., Niikura, C. and Hikita, H. (2009c). Defect states in hydrogenated microcrystalline silicon, *Kotai Butsuri* (Solid State Physics), **44**, pp. 211–225 (in Japanese).

- Morigaki, K. and Ogihara, C. (2006) *Springer Handbook of Electronic and Photonic Materials*, eds. Kasap, S. and Capper, P., "Structural, Optical, and Electronic Properties" (Springer, New York), pp. 565–580.
- Morigaki, K. and Onda, M. (1972). Resistivity decrease due to donor spin resonance in n-type germanium, *J. Phys. Soc. Jpn.*, **33**, pp. 1031–1046.
- Morigaki, K., Sano, Y. and Hirabayashi, I. (1982). Defect creation by optical excitation in hydrogenated amorphous silicon as elucidated by optically detected magnetic resonance, *J. Phys. Soc. Jpn.*, **51**, pp. 147–152.
- Morigaki, K., Takeda, K., Hikita, H., Ogihara, C. and Roca i Cabarrocas, P. (2008). The kinetics of the light-induced defect creation in hydrogenated amorphous silicon—stretched exponential relaxation-, *J. Non-Cryst. Solids*, **354**, pp. 2131–2134.
- Morigaki, K., Takeda, K., Hikita, H. and Roca i Cabarrocas, P. (2005a). Light-induced defect creation in hydrogenated polymorphous silicon, *Mat. Sci. Eng. B*, **121**, pp. 34–41.
- Morigaki, K., Takeda, K., Hikita, H. and Roca i Cabarrocas, P. (2005b). Light-induced defect creation in hydrogenated polymorphous silicon during repeated cycles of illumination and annealing, *Philos. Mag.*, **85**, pp. 3393–3407.
- Morigaki, K., Takeda, K., Hikita, H. and Roca i Cabarrocas, P. (2007). Dispersive processes of light-induced defect creation in hydrogenated amorphous silicon, *Solid State Commun.*, **142**, pp. 232–236.
- Morigaki, K. and Toyotomi, S. (1971). Spin energy transfer from donor spin system to mobile electron system in P-doped Si, *J. Phys. Soc. Jpn.*, **30**, pp. 1207–1208.
- Morigaki, K., Yamaguchi, M., Hirabayashi, I. and Hayasi, R. (1987). *Disordered Semiconductors*, ed. Kastner, M. A., Thomas, G. A. and Ovshinsky, S. R., "Defects in a-Si:H" (Plenum Press, New York) pp. 415–424.
- Morigaki, K. and Yoshida, M. (1985). Gap states in hydrogenated amorphous silicon. The origins of the light-induced ESR and the low-energy luminescence, *Philos. Mag. B*, **52**, pp. 289–298.
- Mott, N. F. (1970). Conduction in non-crystalline systems IV. Anderson localization in a disordered lattices, *Philos. Mag.*, **22**, pp. 7–29.
- Mott, N. F. (1978). Recombination: A survey, *Solid State Electron.*, **21**, pp. 1275–1280.
- Mott, N. F. and Davis, E. A. (1979) *Electronic Processes in Non-Crystalline Materials*, 2nd Ed. (Clarendon Press, Oxford).

- Movaghar, B., Ries, B. and Schweitzer, L. (1980). Theory of the resonant and non-resonant luminescence changes in amorphous silicon, *Philos. Mag. B*, **41**, pp. 141–157.
- Müller, J., Finger, F., Carius, R. and Wagner, H. (1999). Electron spin resonance investigation of electronic states in hydrogenated microcrystalline silicon, *Phys. Rev. B*, **60**, pp. 11666–11677.
- Murayama, K., Morigaki, K. and Kanzaki, H. (1975). Exchange effects in optical detection ESR and dynamics of the optical pumping cycle of F-centers in KCl, *J. Phys. Soc. Jpn.*, **38**, pp. 1623–1633.
- Nakagawa, N., Shimakawa, K., Itoh, T. and Ikeda, Y. (2010). Dynamics of principal photoinduced effects in amorphous chalcogenides: In-situ simultaneous measurements of photodarkening, volume changes, and defect creation, *Phys. Status Solidi C*, **7**, pp. 857–860.
- Nakamura, N., Takahama, T., Isomura, M., Nishikuni, M., Tarui, H., Wakisaka, K., Tsuda, S., Nakano, S., Kishi, Y. and Kuwano, Y. (1992). Irradiation-temperature dependence of the light-induced effect in a-Si solar cells, *Jpn. J. Appl. Phys.*, **31**, pp. 1267–1271.
- Nelson, R. J. (1977). Long-lifetime photoconductivity effect in n-type GaAlAs, *Appl. Phys. Lett.*, **31**, pp. 351–353.
- Neto, A. L. B., Dylla, T., Klein, S., Repmann, T., Lambertz, A., Carius, R. and Finger, F. (2004). Defects and structure of hydrogenated microcrystalline silicon films deposited by different techniques, *J. Non-Cryst. Solids*, **338–340**, pp. 168–172.
- Nickel, N. H., Jackson, W. B. and Johnson, N. M. (1993). Light-induced creation of metastable paramagnetic defects in hydrogenated polycrystalline silicon, *Phys. Rev. Lett.*, **71**, pp. 2733–2736.
- Nielsen, B. Bech, Johannesen, P., Stallinga, P. and Nielsen, K. Bonde (1997). Identification of the silicon vacancy containing a single hydrogen atom by EPR, *Phys. Rev. Lett.*, **79**, pp. 1507–1510.
- Niikura, C., Roca I Cabarrocas, P. and Bourée, J-E. (2011). Structural properties of microcrystalline Si films prepared by hot-wire/catalytic chemical vapor deposition under conditions close to the transition from amorphous to microcrystalline growth, *Thin Solid Films*, **519**, pp. 4502–4505.
- Nishi, Y. (1971). Study of silicon-silicon dioxide structure by electron spin resonance I, *Jpn. J. Appl. Phys.*, **10**, pp. 52–62.
- Nonomura, S., Yoshida, N., Gotoh, T., Sakamoto, T., Kondo, M., Matsuda, A. and Nitta, S. (2000). The light-induced metastable lattice expansion

- in hydrogenated amorphous silicon, *J. Non-Cryst. Solids*, **266–269**, pp. 474–480.
- Ogihara, C. (1998). Frequency resolved spectroscopy in a-Si:H/a-Si<sub>1-x</sub>N<sub>x</sub>:H multilayers and band-edge modulated a-Si<sub>1-x</sub>N<sub>x</sub>:H alloys, *J. Non-Cryst. Solids*, **227–230**, pp. 517–522.
- Ogihara, C., Inagaki, Y. and Morigaki, K. (2011). Light-induced effects on the radiative recombination rate of electron-hole pairs in a-Si:H, *Phys. Status Solidi C*, **8**, pp. 2792–2795.
- Ogihara, C., Inagaki, Y., Taketa, A. and Morigaki, K. (2012). Thermal quenching of defect photoluminescence and recombination rates of electron-hole pairs in a-Si:H, *J. Non-Cryst. Solids*, **358**, pp. 2004–2006.
- Ogihara, C., Nomiya, T., Yamamoto, H., Nakanishi, K., Harada, J., Yu, X. and Morigaki, K. (2006). Light-induced creation of defects related to low energy photoluminescence in hydrogenated amorphous silicon, *J. Non-Cryst. Solids*, **352**, pp. 1064–1067.
- Ogihara, C., Takemura, H., Yoshida, H. and Morigaki, K. (2000). Lifetime distribution of PL under pulsed excitation in hydrogenated amorphous silicon based films, *J. Non-Cryst. Solids*, **266–269**, pp. 574–577.
- Ogihara, C., Yu, X. and Morigaki, K. (2008). Analysis of lifetime distribution of defect luminescence in hydrogenated amorphous silicon, *J. Non-Cryst. Solids*, **354**, pp. 2121–2125.
- Oheda, H. (1999). Real-time modulation of Si-H vibration in hydrogenated amorphous silicon, *Phys. Rev. B*, **60**, pp. 16531–16542.
- Palinginis, K. C., Cohen, J. C., Guha, S. and Yang, J. C. (2001). Experimental evidence indicating a global mechanism for light-induced degradation in hydrogenated amorphous silicon, *Phys. Rev. B*, **63**, pp. 201203-1-201203-4.
- Pankove, J. I. and Berkeyheiser, J. E. (1980). Light-induced radiative recombination centers in hydrogenated amorphous silicon, *Appl. Phys. Lett.*, **37**, pp. 705–706.
- Pantelides, S. T., Chapter 7 “DX Centers in III–V Alloys” (Gordon and Breach Science Publishers, Switzerland), pp. 591–541.
- Pfanner, G., Freysoldt, C. and Neugebauer, J. (2011). *Ab initio* study of electron paramagnetic resonance hyperfine structure of the silicon dangling bond: Role of the local environment, *Phys. Rev. B*, **83**, pp. 144110-1-144110-8.
- Poissant, Y., St’ahel, P. and Roca i Cabarrocas, P. (2000). Effects of temperature on the kinetics of metastable defect creation in polymorphous and

- amorphous silicon materials and solar cells, *Proc. 16<sup>th</sup> European Photovoltaic and Solar energy conference*, pp. 377–380.
- Redfield, D. (1986). Reversibility of recombination-induced defect reactions in amorphous Si:H, *Appl. Phys. Lett.*, **49**, pp. 1517–1518.
- Redfield, D. (1992). Interpretation of stretched-exponential defect kinetics in a-Si:H, *MRS Proc.*, **258**, pp. 341–346.
- Redfield, D. and Bube, R. H. (1990). Identification of defects in amorphous silicon, *Phys. Rev. Lett.*, **65**, pp. 464–467.
- Redfield, D. and Bube, R. H. (1996) *Photoinduced Defects in Semiconductors* (Cambridge University Press, UK).
- Roca i Cabarrocas, P. (2000). Plasma enhanced chemical vapor deposition of amorphous, polymorphous and microcrystalline silicon films, *J. Non-Cryst. Solids*, 266–269, pp. 31–37.
- Roca i Cabarrocas, P., Châbane, N., Kharchenko, A. V. and Tchakarov, S. (2004). Polymorphous silicon thin films produced in dusty plasmas: application to solar cells, *Plasma Phys. Control. Fusion*, **46**, pp. B235–B243.
- Roca i Cabarrocas, P., Fontcuberta i Morral, A. and Possant, Y. (2002). Growth and optoelectronic properties of polymorphous silicon thin films, *Thin Solid Films*, **403–404**, pp. 39–46.
- Roca i Cabarrocas, P., Hamma, S., Sharma, S. N., Viera, G., Bertran, E. and Costa, J. (1989). Nanoparticle formation in low-pressure silane plasmas: bridging the gap between a-Si:H and  $\mu\text{-Si}$  films, *J. Non-Cryst. Solids*, **227–230**, pp. 871–875.
- Roy, D., Longeaud, C., Saadane, O., Gueunier, M. E., Vignoli, S., Butté, R., Meaudre, R. and Meaudre, M. (2002). Evolution with light-soaking of polymorphous material prepared at 423 K, *J. Non-Cryst. Solids*, **299–302**, pp. 482–486.
- Saito, R. and Murayama, K. (1987), A universal distribution function of relaxation in amorphous materials, *Solid State Commun.*, **63**, pp. 625–627.
- Saleh, Z. M., Tarui, H., Ninomiya, K., Takahama, T., Nakashima, Y., Nakamura, N., Haku, H., Wakisaka, K., Tanaka, M., Tsuda, S., Nakano, S., Kishi, M. and Kuwano, Y. (1992). Transient light-induced ESR investigations of the role of hydrogen in the stability of a-Si:H, *Jpn. J. Appl. Phys.*, **31**, pp. 995–998.
- Saleh, Z. M., Tarui, H., Takahama, T., Nakamura, N., Nishikuni, M., Tsuda, S., Nakano, S. and Kuwano, Y. (1993). Trapping and recombination of

- photogenerated carriers in as-grown high-temperature-annealed and light-soaked a-Si:H, *Jpn. J. Appl. Phys.*, **32**, pp. 3376–3384.
- Schmidt, J. and Solomon, I. (1966). Modulation de la photoconductivité dans le silicium á basse temperature par resonance magnétique électronique des impurities peu profondes, *Compt. Rend.*, **263**, pp. 169–172.
- Schultz, N. and Taylor, P. C. (1999). Low temperature kinetics for the growth and decay of band-tail carriers and dangling bonds in hydrogenated amorphous silicon, *MRS Proc.*, **557**, pp. 353–358.
- Schultz, N., Vardeny, Z. V. and Taylor, P. C. (1997). *Spin dependent photoinduced absorption in a-Si:H*, *MRS Proc.*, Vol. 467, eds. Wagner, S., Hack, M., Schiff, E. A., Schropp, R. and Shimizu, I. (*MRS Proc.*, Pittsburgh), pp. 179–183.
- Schumm, G., Jackson, W. B. and Street, R. A. (1993). Nonequilibrium occupancy of tail states and defects in a-Si:H: Implication for defect structure, *Phys. Rev. B*, **48**, pp. 14198–14207.
- Senda, M., Yoshida, N. and Shimakawa, K. (1999). Kinetics of photoinduced defect creation in amorphous semiconductors: analogy to a logistic equation in a biological system, *Philos. Mag. Lett.*, **79**, pp. 375–381.
- Shimakawa, K. (1985). Exciton recombination in amorphous chalcogenides, *Phys. Rev. B*, **31**, pp. 4012–4014.
- Shimakawa, K. and Elliott, S. R. (1988). Reversible photoinduced change of ac conduction in amorphous  $As_2S_3$  films, *Phys. Rev. B*, **38**, pp. 12479–12482.
- Shimakawa, K., Ikeda, Y. and Kugler, S. (2004). *Non-Crystalline Materials for Optoelectronics* Vol. 1, ed. Popescu, M., Chapter 5 “Fundamental Optoelectronic Processes in Amorphous Chalcogenides” (INOE Publishing House, Bucharest), pp. 103–130.
- Shimakawa, K., Inami, S. and Elliott, S. R. (1990). Reversible photoinduced change of photoconductivity in amorphous chalcogenide films, *Phys. Rev. B*, **42**, pp. 11857–11861.
- Shimakawa, K., Kolobov, A. V. and Elliott, S. R. (1995). Photoinduced effects and metastability in amorphous semiconductors and insulators, *Adv. Phys.*, **44**, pp. 475–588.
- Shimakawa, K., Meherun-Nessa, Ishida, H. and Ganjoo, A. (2004). Quantum efficiency of light-induced defect creation in hydrogenated amorphous silicon and amorphous  $As_2Se_3$ , *Philos. Mag. Lett.*, **84**, pp. 81–89.
- Shimizu, T. (2004). Stabler-Wronski effect in hydrogenated amorphous silicon and related alloy films, *Jpn. J. Appl. Phys.*, **43**, pp. 3257–3268.

- Shinozuka, Y. and Toyozawa, Y. (1979). Self-trapping in mixed crystal-Clustering, dimensionality, percolation, *J. Phys. Soc. Jpn.*, **46**, pp. 505–514.
- Singh, J. and Shimakawa, K. (2003) *Advances in Amorphous Semiconductors* (Taylor & Francis, London).
- Solomon, I. (1972) *Spin-dependent transport in semiconductors*, Proc. 11th International Conf. Phys. Semiconductors, Vol. 1 (Polish Sci. Pub., Warsaw), pp. 27–37.
- Solomon, I. (1976). Spin-dependent recombination in a silicon p-n junction, *Solid State Commun.*, **20**, pp. 215–217.
- Solomon, I. (1979) *Amorphous Semiconductors*, ed. Brodsky, M. H., Chapter 7, “Spin effects in Amorphous Semiconductors” (Springer, Berlin), pp. 189–213.
- Solomon, I. and Bhatnagar, M. (1992). Transport dependant du spin dans une diode cristalline, *Compt. Rend*, **314**, pp. 1133–1138.
- Solomon, I., Biegelsen, D. and Knights, J. C. (1977). Spin-dependent photoconductivity in n-type and p-type amorphous silicon, *Solid State Commun.*, **22**, pp. 505–508.
- Stachowitz, R., Schubert, M. and Fuhs, W. (1994). Frequency-resolved spectroscopy and its application to low-temperature geminate recombination in a-Si:H, *Philos. Mag. B*, **70**, pp. 1219–1230.
- Staabler, D. L. and Wronski, C. R. (1977). Reversible conductivity changes in discharge-produced amorphous Si, *Appl. Phys. Lett.*, **31**, pp. 292–294.
- Staabler, D. L. and Wronski, C. R. (1980). Optically induced conductivity changes in discharge-produced hydrogenated amorphous silicon, *J. Appl. Phys.*, **51**, pp. 3262–3268.
- Stradins, P. and Fritzsche, H. (1996). Light-induced metastable changes in defect density and photoconductivity of a-Si:H between 4.2 and 300 K, *J. Non-Cryst. Solids*, **198–200**, pp. 432–435.
- Stradins, P., Kondo, M. and Matsuda, A. (2000). A study of light-induced degradation of a-Si:H by nanosecond laser pulse pairs of variable delay, *J. Non-Cryst. Solids*, **266–269**, pp. 405–409.
- Street, R. A. (1991). *Hydrogenated Amorphous Silicon* (Cambridge University Press, UK).
- Street, R. A. and Biegelsen, D. K. (1980). Luminescence and ESR studies of defects in hydrogenated amorphous silicon, *Solid State Commun.*, **33**, pp. 1159–1162.

- Street, R. A., Knights, J. C. and Biegelsen, D. K. (1978). Luminescence studies of plasma-deposited hydrogenated silicon, *Phys. Rev. B*, **18**, pp. 1880–1891.
- Street, R. A. and Mott, N. F. (1975). States in the gap in glassy semiconductors, *Phys. Rev. Lett.*, **35**, pp. 1293–1296.
- Stutzmann, M. and Biegelsen, D. K. (1983). Electron-spin-lattice relaxation in amorphous silicon and germanium, *Phys. Rev. B*, **28**, pp. 6256–6261.
- Stutzmann, M. and Biegelsen, D. K. (1986). Electron-nuclear double-resonance experiments in hydrogenated amorphous silicon, *Phys. Rev. B*, **34**, pp. 3093–3107.
- Stutzmann, M., Brandt, M. S. and Bayerl, M. W. (2000). Spin-dependent processes in amorphous and microcrystalline silicon: a survey, *J. Non-Cryst. Solids*, **266–269**, pp. 1–22.
- Stutzmann, M., Jackson, W. B. and Tsai, C. C. (1985). Light-induced metastable defects in hydrogenated amorphous silicon: A systematic study, *Phys. Rev. B*, **32**, pp. 23–47.
- Stutzmann, M., Rossi, M. C. and Brandt, M. S. (1994). Pulsed-light soaking of hydrogenated amorphous silicon, *Phys. Rev. B*, **50**, pp. 11592–11605.
- Su, T., Ju, T., Yan, B., Yang, J., Guha, S. and Taylor, P. C. (2008). ESR study of the hydrogenated nanocrystalline silicon thin films, *J. Non-Cryst. Solids*, **354**, pp. 2231–2234.
- Su, T., Taylor, P. C., Ganguly, G. and Carlson, D. E. (2002). Direct role of hydrogen in the Staebler-Wronski effect in hydrogenated amorphous silicon, *Phys. Rev. Lett.*, **89**, pp. 015502-1–015502-4.
- Su, T., Taylor, P. C., Ganguly, G. and Carlson, D. E. (2004). A hydrogen-related defect and the Staebler-Wronski effect in hydrogenated amorphous silicon, *J. Non-Cryst. Solids*, **338–340**, pp. 357–360.
- Sumi, H. (1984). Dynamic defect reaction induced by multiphonon nonradiative recombination of injected carriers at deep levels in semiconductors, *Phys. Rev. B*, **29**, pp. 4616–4630.
- Sumi, H. (1988). Nonradiative electron-hole recombination at deep-level defects and recombination enhanced defect reactions in semiconductors, *Kotai Butsuri (Solid State Physics)*, **23**, pp. 221–230 (in Japanese).
- Takeda, K., Hikita, H., Kimura, Y., Yokomichi, H., Yamaguchi, M. and Morigaki, K. (1997). Light-induced annealing of dangling bonds in a-Si:H, *Jpn. J. Appl. Phys.*, **36**, pp. 991–996.
- Takeda, K., Hikita, H., Kimura, Y., Yokomichi, H. and Morigaki, K. (1998). Electron spin resonance study of light-induced annealing of



- dangling bonds in glow discharge hydrogenated amorphous silicon: Deconvolution of electron spin resonance spectra, *Jpn. J. Appl. Phys.*, **37**, pp. 6309–6317.
- Takeda, K., Hikita, H., Kimura, Y., Yokomichi, H., Izumi, T. and Morigaki, K. (2000). Electron spin resonance spectra and their thermal annealing effects in hydrogenated amorphous silicon, *Proceedings of the School of Engineering, Tokai Univ., Series E*, **25**, 17–25.
- Takeda, K., Morigaki, K., Hikita, H. and Roca i Cabarrocas P. (2008). Thermal annealing effects of dangling bonds in hydrogenated polymorphous silicon, *J. Appl. Phys.*, **104**, pp. 053715-1-053715-6.
- Takemura, H., Ogihara, C. and Morigaki, K. (2002). Lifetime resolved study of photoluminescence under pulsed excitation in a-Si:H based films, *J. Phys. Soc. Jpn.*, **71**, pp. 625–629.
- Takenaka, H., Ogihara, C. and Morigaki, K. (1988). Time-resolved optically detected magnetic resonance experiment on conduction band tail electrons and A centre in hydrogenated amorphous silicon, *J. Phys. Soc. Jpn.*, **57**, pp. 3858–3867.
- Tanaka, Ke. and Shimakawa, K. (2011) *Amorphous Chalcogenide Semiconductors and Related Materials* (Springer, New York).
- Taylor, P. C. (2006). The localization of electrons in amorphous semiconductors: A twenty-first century perspective, *J. Non-Cryst. Solids*, **352**, pp. 839–850.
- Tiedje, T. (1984) *Semiconductors and Semimetals 21 Part C*, ed. Pankove, J. I., Chapter 6 “Information about band-tail states from time-of-flight experiments” (Academic Press, Orlando), pp. 207–238.
- Timilsina, R. and Biswas, P. (2010). Theoretical study of hydrogen microstructure in models of hydrogenated amorphous silicon, *Phys. Status Solidi A*, **207**, pp. 609–612.
- Toyoda, Y. and Hayashi, Y. (1970). Bolometric detection of ESR in P-doped Si at low temperature, *J. Phys. Soc. Jpn.*, **29**, pp. 247–248.
- Toyoda, Y. and Hayashi, Y. (1971). The spin dependent conductivity in P-doped Si, *J. Phys. Soc. Jpn.*, **30**, 1511–1512.
- Toyotomi, S. (1974). Resistance change induced by electron spin resonance and spin relaxation in phosphorus-doped silicon, *J. Phys. Soc. Jpn.*, **37**, pp. 130–139.
- Toyotomi, S. and Morigaki, K. (1970). Microwave hot electron effect and resistivity decrease due to donor spin resonance in P-doped Si, *Solid State Commun.*, **8**, pp. 1307–1308.

- Toyozawa, Y. (2003) *Optical Processes in Solids* (Cambridge University Press, UK).
- Tsang, T. and Street, R. A. (1979). Recombination in plasma-deposited amorphous Si:H. Luminescence decay, *Phys Rev. B*, **19**, pp. 3027–3040.
- Tzanetakos, P., Kopidakis, N., Androulidaki, M., Kalouzos, C., Stradins, P. and Fritzsche, H. (1996). Short-laser-pulse and steady-light induced degradation of intrinsic, p-type and compensated a-Si:H, *J. Non-Cryst. Solids*, **198–200**, pp. 458–461.
- Uchino, T., Takahashi, M. and Yoko, T. (2001).  $E'$  centers in amorphous SiO<sub>2</sub> revisited: A new look at an old problem, *Phys. Rev. Lett.*, **86**, pp. 5522–5525.
- Uchino, T. and Yoko, T. (2003). Mechanism of electron localization at edge-sharing units in amorphous SiO<sub>2</sub>, *Phys. Rev. B*, **68**, pp. 041201- –041201-4.
- Umeda, T., Yamasaki, S., Isoya, J., Matsuda, A. and Tanaka, K. (1996). Electronic structure of band-tail electrons in a-Si:H, *Phys. Rev. Lett.*, **77**, pp. 4600–4603.
- Vanderbilt, D. and Joannopoulos, J. D. (1980). Theory of defect states in glassy selenium, *Phys. Rev. B*, **22**, pp. 2927–2939.
- Vanderbilt, D. and Joannopoulos, J. D. (1981). Theory of defect states in glassy As<sub>2</sub>Se<sub>3</sub>, *Phys. Rev. B*, **23**, pp. 2596–2606.
- Vardeny, Z. and Olszakier, M. (1987). Infrared photomodulation spectroscopy of band tail states in a-Si:H and a-Si:F, *J. Non-Cryst. Solids*, **97–98**, pp. 109–112.
- Vetterl, O., Finger, F., Carius, R., Hapke, P., Houben, L., Kluth, O., Lambertz, A., Mück, A., Rech, B. and Wagner, H. (2000). Intrinsic microcrystalline silicon: A new material for photovoltaics, *Sol. Energy Mat. Sol. Cells*, **62**, pp. 97–108.
- Vignoli, S., Meaudre, R. and Meaudre, M. (1996). Metastable defect creation and annealing under illumination in intrinsic hydrogenated amorphous silicon deposited from helium–silane mixtures, *Philos. Mag. B*, **73**, pp. 261–276.
- Watkins, G. D. and Corbett, J. W. (1964). Defects in irradiated silicon: Electron paramagnetic resonance and electron–nuclear double resonance of the Si-E center, *Phys. Rev.*, **134**, A1359–A1377.
- Watkins, G. D. (1965). *A review of EPR studies in irradiated silicon*, *Proc. Radiation Damage in Semiconductors* (Dunod, Paris), pp. 97–113.

- Watkins, G. D. (2000). *Handbook of Semiconducto Technology*, Vol. 1, eds. Jackson, K. A. and Schröter, W., Chapter 3 "Intrinsic Point Defectsin Semiconductors 1999" (Wiley-VCH, Weinheim), pp. 121–165.
- Weeks, J. D., Tully, J. C. and Kimerling, L. C. (1975). Theory of recombination-enhanced defect reactions in semiconductors. *Phys. Rev. B*, **12**, pp. 3286–3292.
- Wilson, D. K. and Feher, G. (1961). Electron spin resonance experiments on donors in silicon III. Investigation of excited states by the application of uniaxial stress and their importance in relaxation processes, *Phys. Rev.*, **124**, pp. 1068–1084.
- Winer, K., Hirabayashi, I. and Ley, L. (1988a). Exponential conduction-band tail in P-doped a-Si:H, *Phys. Rev. Lett.*, **60**, 2697–2700.
- Winer, K., Hirabayashi, I. and Ley, L. (1988b). Distribution of occupied near-surface band-gap states in a-Si:H, *Phys. Rev. B*, **38**, pp. 7680–7693.
- Wraback, M. and Tauc, J. (1992). Direct measurement of the hot carrier cooling rate in a-Si:H using femtosecond 4 eV pulses, *Phys. Rev. Lett.*, **69**, pp. 3682–3685.
- Yamaguchi, M. and Morigaki, K. (1993). Optical properties of amorphous-microcrystalline mixed-phase Si:H films, *J. Phys. Soc. Jpn.*, **62**, pp. 2915–2923.
- Yamaguchi, M. and Morigaki, K. (1997). Photoluminescence and optically detected magnetic resonance in a-Si:H/a-Si<sub>3</sub>N<sub>4</sub>:H multilayers, *Phys. Rev. B*, **55**, pp. 2368–2377.
- Yamaguchi, M. and Morigaki, K. (1999). Effect of hydrogen dilution on the optical properties of hydrogenated amorphous silicon prepared by plasma deposition, *Philos. Mag. B*, **79**, pp. 387–405.
- Yamaguchi, M., Morigaki, K. and Nitta, S. (1989). Observation of light-induced phenomena by photoluminescence and optically detected magnetic resonance in a-Si:H, *J. Phys. Soc. Jpn.*, **58**, pp. 3828–3841.
- Yamaguchi, M., Morigaki, K. and Nitta, S. (1991). Observation of light-induced phenomena by photoluminescence and optically detected magnetic resonance in a-Si<sub>1-x</sub>N<sub>x</sub>:H, *J. Phys. Soc. Jpn.*, **60**, pp. 1769–1791.
- Yamasaki, S. and Isoya, J. (1993). Pulsed-ESR study of light-induced metastable defect in a-Si:H, *J. Non-Cryst. Solids*, **164–166**, pp. 169–174.
- Yamasaki, S., Okushi, H., Matsuda, A. and Tanaka, K. (1990). Origin of optically induced electron-spin resonance in hydrogenated amorphous silicon, *Phys. Rev. Lett.*, **65**, pp. 756–759.

- Yan, B, Schultz, N. A., Efros, A. L. and Taylor, P. C. (2000). Universal distribution of residual carriers in tetrahedrally coordinated amorphous semiconductors, *Phys. Rev. Lett.*, **84**, pp. 4180–4183.
- Yokomichi, H., Hirabayashi, I. and Morigaki, K. (1987). Electron-nuclear double resonance of dangling-bond centres in a-Si:H, *Solid State Commun.*, **61**, pp. 697–701.
- Yokomichi, H. and Morigaki, K. (1987). Electron-nuclear double resonance of dangling bonds centres associated with hydrogen incorporation in a-Si:H, *Solid State Commun.*, **63**, pp. 629–632.
- Yokomichi, H. and Morigaki, K. (1991). The nature of structural defects and thermal equilibrium effects on doped a-Si:H as elucidated by electron nuclear double resonance measurements, *J. Non-Cryst. Solids*, **137–138**, pp. 183–186.
- Yokomichi, H. and Morigaki, K. (1993). Clustered dangling bonds in a-Si:H as elucidated by ENDOR measurements, *Solid State Commun.*, **85**, pp. 759–761.
- Yokomichi, H. and Morigaki, K. (1996). Evidence for existence of hydrogen-related dangling bonds in hydrogenated amorphous silicon, *Philos. Mag. Lett.*, **73**, pp. 283–287.
- Yonezawa, F., Sakamoto, S. and Hori, M. (1991). Ab-initio molecular dynamics simulations of amorphous silicon, *J. Non-Cryst. Solids*, **137–138**, pp. 135–140.
- Yoshida, M. and Morigaki, K. (1989). Triplet exciton states in a-Si:H and its alloys as elucidated by optically detected magnetic resonance measurements, *J. Phys. Soc. Jpn.*, **58**, pp. 3371–3382.
- Yoshida, M. and Taylor, P. C. (1992). Correlations of changes in ESR and PL with light soaking in a-Si:H, *MRS Proc.*, **258**, pp. 347–352.
- Yoshida, M., Yamaguchi, M. and Morigaki, K. (1989). Nature of triplet excitons in a-Si:H, its alloys and a-Si:H multilayers as elucidated by optically detected magnetic resonance measurements, *J. Non-Cryst. Solids*, **114**, pp. 319–321.
- Zafar, S. and Schiff, E. A. (1989). Hydrogen-mediated model for defect metastability in hydrogenated amorphous silicon, *Phys. Rev. B*, **40**, 5235–5238.
- Zhang, S. B. and Branz, H. M. (2000). Nonradiative electron-hole recombination by a low-barrier pathway in hydrogenated silicon semiconductors, *Phys. Rev. Lett.*, **84**, pp. 967–970.

- Zhang, Q., Nishino, T., Kumeda, M. and Shimizu, T. (1995). Light-induced annealing of photocreated dangling bonds in hydrogenated amorphous silicon, *Jpn. J. Appl. Phys.*, **34**, pp. 483–486.
- Zhou, J. H., Kumeda, M. and Shimizu, T. (1995). Light-induced ESR in variously treated hydrogenated amorphous silicon, *Jpn. J. Appl. Phys.*, **34**, pp. 3982–3986.



# Index

- a-As<sub>2</sub>S<sub>3</sub> *see* amorphous As<sub>2</sub>S<sub>3</sub>
- a-As<sub>2</sub>Se<sub>3</sub> *see* amorphous As<sub>2</sub>Se<sub>3</sub>
- a-Se *see* amorphous selenium
- ab initio* molecular dynamics
  - computer simulation 41
- a-Ch *see* amorphous chalcogenide
- amorphous network 25–28, 52, 54, 85, 98, 99, 117, 128, 129, 136–138
- amorphous As<sub>2</sub>S<sub>3</sub> (a-As<sub>2</sub>S<sub>3</sub>) 163–166, 168
- amorphous As<sub>2</sub>Se<sub>3</sub> (a-As<sub>2</sub>Se<sub>3</sub>) 163, 164, 166, 169
- amorphous selenium (a-Se) 164, 165
  - hydrogenated (a-Si:H) 2, 3, 5, 7, 22–31, 35–38, 40–42, 44–79, 81–141, 153–164,
- amorphous chalcogenide (a-Ch) 2, 3, 91, 161, 164, 166
- anisotropic hyperfine interaction
  - constant 50, 134
- anisotropic magnetic center 145–147
- antibonding orbital 35, 37
  
- band tail 28, 35, 37, 152, 155, 158
- biexciton 85
  
- c-Si *see* crystalline silicon
- capture cross section 16
- characteristic time 83, 99, 170
- charged Si-dangling bond 26
- clustered phase 87
  
- column boundary 145
- computer simulation 41, 164
- conduction band edge 8
- conduction band tail state 155
- conduction electron ESR 149
- configuration coordinate model 13, 17, 19, 65, 66
- constant photocurrent method (CPM) 69
- continuous random network model 26
- coordination number 29, 166, 167
- crystalline phase 155
- crystalline silicon (c-Si) 2, 5
- crystalline volume fraction 26, 51, 52, 149
- crystallite 27, 148
- crystallite fraction 27
  
- dangling bond (DB) 3, 21–23, 25, 26, 29, 34–40, 48, 49, 51–53, 57–59, 67, 69–72, 75–140, 144, 148, 151, 155, 157, 167
- deep center 30, 33
- deep gap state 54
- deep level 13, 20
- deep tail state 64
- defect 10, 11, 13–15, 17, 22–24, 57–59, 64, 65, 67, 70–75, 77–79, 81–87, 89–91, 99–105, 107–109, 111–113, 115–121
- defect creation efficiency 74, 82, 123
- defect PL 55, 57–59, 64, 65, 67, 70
- delocalized state 27

- diffusion 10–12, 86, 104, 116, 117, 135, 138
- dilute phase 74, 75, 87
- disordered system 25, 100
- dispersive hydrogen diffusion 104, 135
- dispersive parameter 83
- distant electron–hole pair 4, 32
- distant ENDOR 108, 109
- DX center 17, 20
- electronegativity 41, 71
- electron–hole pair 4, 32, 60–64, 85
- electron–nuclear double resonance (ENDOR) 2, 3, 28, 29, 5, 36, 40, 43, 45, 48, 56, 108–112, 164
- electron–phonon coupling 56, 65, 66
- electron spin resonance (ESR) 2–5, 20, 28–30, 35, 36, 48, 49, 67, 71–73, 101, 104–106, 108, 110, 111, 116, 117, 122–129, 132–134, 137, 143–154, 163, 164
- ENDOR *see* electron–nuclear double resonance
- ESR *see* electron spin resonance
- exchange interaction 47
- exciton 168
- excitonic PL 62
- extrinsic self-trapping 41
- Fermi-type contact interaction 50
- fill factor 118, 119
- fourfold-coordinated Si 20
- fractal dimension 140
- fractal-time random walk 135, 140
- fractal-time random walk model 135
- frequency-resolved spectroscopy (FRS) 59, 60, 62, 63
- FWHM 44, 46, 48, 55, 155–157
- Ga<sub>1-x</sub>Al<sub>x</sub>As 17, 18, 20
- gap state 29, 54, 56, 57, 59, 65, 68, 69
- GaAs 12, 14, 18, 20, 21
- gas dilution ratio 143
- Gaussian spin-packet 107, 144
- geminate pair 61
- geminate recombination 61
- glass transition temperature T<sub>g</sub> 162
- Godet model 97, 98
- grain boundary 21, 158
- g-shift 37–39
- g-tensor 38
- g-value 2, 37, 39, 46, 47, 136, 144, 146–149, 157
- H-DB 38, 40, 85
- hot-wire chemical vapor deposition (hot-wire CVD) 26, 39
- H-related dangling bond 104, 116
- hydrogen bonding scheme 137
- hydrogenated microcrystalline silicon ( $\mu\text{c-Si:H}$ ) 2, 3, 26, 30, 35, 38–40, 143–147, 149–151, 153–160
- hydrogenated polycrystalline silicon (poly-Si:H)
- hydrogenated polymorphous silicon (pm-Si:H) 26, 39
- hydrogen-collision model 102
- hydrogen-mediated model 75, 87, 88
- hydrogen-related dangling bond 40, 48, 49, 51, 78, 79, 99, 104,



- 108, 111, 112, 114, 131, 133, 134
- hydrogen-site energy 136
- hyperfine interaction constant 50, 51, 108, 134
- hyperfine structure 49, 108
- impurity-involved model 75, 90
- isotropic hyperfine interaction constant 50, 51
- k-selection rule 54
- LESR *see* light-induced electron spin resonance
- lifetime distribution 59–63, 99, 136, 138
- light-induced annealing 98, 128–130
- light-induced dangling bond 83, 85, 99, 117–119, 128, 135
- light-induced defect 22–24, 71, 73–75, 77–79, 81–83, 89–91, 97, 99–101, 103–105, 111–113, 117–119, 121, 123, 127, 135–137
- light-induced degradation 137, 138, 159
- light-induced effect 159, 166
- light-induced electron spin resonance (LESR) 30–37, 39, 150, 151, 153, 154, 164, 166
- light-induced hydrogen pair 112
- light-induced structural change 134
- light soaking 34, 35, 42, 117
- local ENDOR 48, 108, 109
- luminescence fatigue 72, 166
- magnetic resonance 28, 54, 155, 157, 164
- matrix ENDOR 45
- metastable hydrogen atom 78, 79, 112
- microcrystalline grain 26
- microcrystalline grain size 26
- microcrystalline phase 39, 51, 52
- microvoid 89
- mid-gap absorption 163, 166
- mobility edge 27
- mobility gap 28
- modulated photo current method (MPC) 101, 102
- monomolecular recombination 82, 95, 122
- MPC *see* modulated photo current method
- multiphonon emission 14, 66
- negative correlation energy  $U$  33
- negative- $U$  center 33, 89
- negatively charged threefold-coordinated Si atom 29, 89
- nongeminate electron–hole pair 4
- nongeminate recombination 61
- nonradiative capture 13
- nonradiative center 33, 65, 148
- nonradiative electron–hole recombination 75, 85
- nonradiative recombination 29, 34, 56–59, 63–67, 69, 70, 72, 129, 134, 158, 166
- nonradiative recombination center 72
- nonradiative recombination rate 57, 65, 70

- normal dangling bond 40, 49, 78, 79, 99, 104, 112, 114, 116, 131
- ODENDOR *see* optically detected ENDOR
- ODMR *see* optically detected magnetic resonance
- optical absorption 28, 29, 67, 69, 101, 121, 162, 168
- optical absorption coefficient 101, 168
- optical gap energy 51–53
- optically detected ENDOR (ODENDOR) 33, 40, 43, 44, 56, 164
- optically detected magnetic resonance (ODMR) 3–5, 28, 33, 34, 36, 37, 40, 45, 46, 48, 54, 56, 57, 62, 117, 143, 155, 157–159, 164
- Pb center 39, 40, 145
- PDS *see* photo-thermal deflection spectroscopy
- PECVD *see* plasma-enhanced chemical vapor deposition
- persistent photoconductivity 16, 18
- photoconductivity 16, 18, 68, 101, 102, 121, 122, 143
- photodarkening 162, 163
- photoemission spectroscopy 28
- photoinduced absorption (PA) 5,
- photoluminescence (PL) 54, 69, 70, 155, 157, 159
- photomodulation-IR absorption spectrum 43, 45, 56
- photo-thermal deflection spectroscopy (PDS) 69
- plasma-enhanced chemical vapor deposition (PECVD) 26, 39
- plasma-enhanced CVD *see* plasma-enhanced chemical vapor deposition
- PL *see* photoluminescence
- platelets 23, 136
- polycrystalline silicon (poly-Si) 21
- positive correlation energy 89, 148
- principal PL 30, 48, 55–59, 64, 65, 67, 69, 70, 158
- pulsed ENDOR technique 110
- QFRS *see* quadrature-frequency resolved spectroscopy
- quadrature-frequency resolved spectroscopy (QFRS) 60
- radiation-induced defect 8, 12,
- radiative recombination 32, 48, 57, 60, 64, 66, 70, 158
- radiative recombination rate 60, 64
- recombination process 65
- repeated LS-annealing cycling 138
- saturated dangling bond density 83, 94, 96, 124, 125
- self-trapped hole 29, 37, 43, 47, 48, 123
- self-trapped hole center 29
- shallow level 20, 151
- shallow tail state 64, 65
- Si-H bond 129
- singlet exciton 62
- solar cell 1, 5, 25, 67, 100, 101, 118, 119
- spatial fluctuation 27
- s-p hybridized orbital 165

- specific weak Si–Si bond 44, 47, 77  
 spin density 22, 23, 77, 118, 120,  
 122, 136, 150–152, 154, 165,  
 166  
 spin-lattice relaxation time 115,  
 153  
 stretched exponential function 32,  
 33, 82, 83, 85, 91–95, 97, 99,  
 101, 135, 136, 140, 170  
 stretched exponential relaxation  
 83, 91, 97, 100  
 stretching vibration 41, 42  
 strong electron–phonon coupling  
 65, 66  
 structural change 41, 134  
 structural defect 28, 29, 40  
 structural factor R 74  
 subbandgap absorption 119  
 subgap illumination 123, 162
- tail electron 35, 37, 45, 47, 48, 151,  
 152  
 tail hole 37, 152  
 tail state 57  
 Te-doped  $\text{Ga}_{1-x}\text{Al}_x\text{As}$  16  
 tetrahedral angle 146  
 tetrahedral coordination 27  
 thermal activation 10, 14, 15  
 thermal activation energy 14, 15  
 thermal annealing 103, 127, 128,  
 135, 136, 139, 140, 164, 166
- thermal stability 164  
 thin film 69  
 thin-film transistor 25  
 threefold-coordinated dopant 90  
 threefold-coordinated Si 25  
 time-resolved ODMR 33, 37, 45, 48  
 transport 28  
 trapping process 79  
 triplet exciton 62  
 triplet state 4, 62  
 tunneling 57  
 type I LESR center 166  
 type II LESR center 166
- vacancy–hydrogen complex 38, 53  
 valence band tail hole 152  
 void 130, 133, 134  
 volume fraction 26, 51, 52, 98, 149
- weak bond 43, 79  
 weak electron–phonon coupling  
 66  
 weak Si–Si bond 32, 34, 35, 37,  
 41–45, 47, 71, 72, 75–79, 87,  
 88, 114, 123, 125  
 weak Si–Si bond breaking model  
 75  
 wrong bond 166

