

## Bibliography

- Adachi, S. (1999). *Optical Constants of Crystalline and Amorphous Semiconductors* (Kluwer Academic, Boston, MA).
- Adams, A. R. (1986). Band-structure engineering for low-threshold high-efficiency semiconductor lasers, *Electron. Lett.*, **22**, 249.
- Akahane, Y., Asano, T., Song, B., and Noda, S. (2003). Investigation of high-Q channel drop filters using donor-type defects in two-dimensional photonic crystal slabs, *Appl. Phys. Lett.*, **83**, 1512.
- Akahane, Y., Asano, T., Song, B., and Noda, S. (2005). Fine-tuned high-Q photonic-crystal nanocavity, *Opt. Express*, **13**, 1202.
- Allison, G., Spasov, S., Patanè, A., Eaves, L., Kozlova, N., Freudenberger, J., Hopkinson, M., and Hill, G. (2008). Electron effective mass and Si-donor binding energy in  $\text{GaAs}_{1-x}\text{N}_x$  probed by a high magnetic field, *Phys. Rev. B*, **77**, 125210.
- Alt, H. C., Egorov, A. Y., Riechert, H., Wiedemann, B., Meyer, J. D., Michelmann, R. W., and Bethge, K. (2000). Infrared absorption study of nitrogen in N-implanted GaAs and epitaxially grown  $\text{GaAs}_{1-x}\text{N}_x$  layers, *Appl. Phys. Lett.*, **77**, 3331.
- Alt, H. C., Egorov, A. Y., Riechert, H., Wiedemann, B., Meyer, J. D., Michelmann, R. W., and Bethge, K. (2001). Local vibrational mode absorption of nitrogen in GaAsN and InGaAsN layers grown by molecular beam epitaxy, *Physica B*, **302–303**, 282.
- Alt, H. C., Messerer, P., Köhler, K., and Riechert, H. (2009). H- and D-related mid-infrared absorption bands in  $\text{Ga}_{1-y}\text{In}_y\text{As}_{1-x}\text{N}_x$  epitaxial layers, *Phys. Status Solidi B*, **246**, 200.
- Altug, H., Englund, D., and Vuckovic, J. (2006). Ultrafast photonic crystal nanocavity laser, *Nat. Phys.*, **2**, 484.
- Alvarez, J., Kleider, J.-P., Trotta, R., Polimeni, A., Capizzi, M., Martelli, F., Mariucci, L., and Rubini, S. (2011). Giant and reversible enhancement of the electrical resistance of  $\text{GaAs}_{1-x}\text{N}_x$  by hydrogen irradiation, *Phys. Rev. B*, **84**, 085331.

- Amidani, L., Ciatto, G., Bondino, F., and Boscherini, F. (2014). Local electronic structure of dilute elements in alloys, *Phys. Rev. B* (submitted).
- Amore Bonapasta, A. (1991). Hydrogen passivation of carbon-doped gallium arsenide, *Phys. Rev. B*, **48**, 8771.
- Amore Bonapasta, A., and Capizzi, M. (1998). Hydrogen as a deep impurity in semiconductors and its interaction with deep centers in III-V compounds, *Defect Diffus. Forum*, **157-159**, 133.
- Amore Bonapasta, A., and Filippone, F. (2003). Local and lattice relaxations in hydrogenated  $\text{GaAs}_{y}\text{N}_{1-y}$  alloys, *Phys. Rev. B*, **68**, 073202.
- Amore Bonapasta, A., Filippone, F., and Giannozzi, P. (2003). Nitrogen passivation by atomic hydrogen in  $\text{GaAs}_{y}\text{N}_{1-y}$  and  $\text{In}_x\text{Ga}_{1-x}\text{As}_y\text{N}_{1-y}$  alloys, *Phys. Rev. B*, **68**, 115202.
- Amore Bonapasta, A., Filippone, F., and Giannozzi, P. (2004). Structure, electronic properties, and formation mechanisms of hydrogen-nitrogen complexes in  $\text{GaP}_y\text{N}_{1-y}$  alloys, *Phys. Rev. B*, **69**, 115207.
- Amore Bonapasta, A., Filippone, F., and Giannozzi, P. (2005). Electron correlation effects on the hydrogen passivation of  $\text{Mn}_x\text{Ga}_{1-x}\text{As}$  dilute magnetic semiconductors, *Phys. Rev. B*, **72**, 121202(R).
- Amore Bonapasta, A., Filippone, F., Giannozzi, P., Capizzi, M., and Polimeni, A. (2002). Structure and passivation effects of mono- and dihydrogen complexes in  $\text{GaAs}_{1-y}\text{N}_y$  alloys, *Phys. Rev. Lett.*, **89**, 216401.
- Amore Bonapasta, A., Filippone, F., and Mattioli, G. (2007). H-induced dangling bonds in H-isoelectronic-impurity complexes formed in  $\text{GaAs}_{1-y}\text{N}_y$  alloys, *Phys. Rev. Lett.*, **98**, 206403.
- Amore Bonapasta, A., and Giannozzi, P. (2000). Effects of strain and local charge on the formation of deep defects in III-V ternary alloys, *Phys. Rev. Lett.*, **84**, 3923.
- Amore Bonapasta, A., and Pavesi, L. (1996). Hydrogen interaction with shallow and deep centers in GaAs, *Int. J. Quantum Chem.*, **57**, 823.
- Amsel, G., and Davies, J. (1969). La microanalyse de l'azote par l'observation directe de réactions nucléaires applications, *Rev. Phys. Appl.*, **4**, 383.
- Andreev, A. D., and O'Reilly, E. P. (2004). Theoretical study of Auger recombination in a GaInNAs 1.3  $\mu\text{m}$  quantum well laser structure, *Appl. Phys. Lett.*, **84**, 1826.
- Anisimov, V. I., Aryasetiawan, F., and Lichtenstein, A. I. (1997). First-principles calculations of the electronic structure and spectra of strongly correlated systems: the LDA+U method, *J. Phys.: Condens. Matter*, **9**, 767.

- Ankudinov, A. L., Ravel, B., Rehr, J. J., and Conradson, S. D. (1998). Real-space multiple-scattering calculation and interpretation of x-ray-absorption near-edge structure, *Phys. Rev. B*, **58**, 7565.
- Ashcroft, N. W., and Mermin, N. D. (1976). *Solid State Physics* (Saunders College/Harcourt Brace College).
- Atlasov, K. A., Calic, M., Karlsson, K. F., Gallo, P., Rudra, A., Dwir, B. and Kapon, E. (2009). Photonic-crystal microcavity laser with site-controlled quantum-wire active medium, *Opt. Express*, **17**, 18178.
- Atlasov, K. A., Felici, M., Karlsson, K. F., Gallo, P., Rudra, A., Dwir, B. and Kapon, E. (2010). 1D photonic band formation in finite-size photonic-crystal waveguides, *Opt. Express*, **18**, 117.
- Atlasov, K. A., Karlsson, K. F., Deichsel, E., Rudra, A., Dwir, B., and Kapon, E. (2007). Site-controlled single quantum wire integrated into a photonic-crystal membrane microcavity, *Appl. Phys. Lett.*, **90**, 153107.
- Atlasov, K. A., Karlsson, K. F., Rudra, A., Dwir, B., and Kapon, E. (2008). Wavelength and loss splitting in directly coupled photonic-crystal defect microcavities, *Opt. Express*, **16**, 16255.
- Atlasov, K. A., Rudra, A., Dwir, B., and Kapon, E. (2011). Large mode splitting and lasing in optimally coupled photonic-crystal microcavities, *Opt. Express*, **19**, 2619.
- Averbeck, R., Jaschke, G., Geelhaar, L., and Riechert, H. (2004). Record-low thresholds of InGaAsN/GaAs SQWs lasers, in *Proc. 19th Int. Semicond. Laser Conf.* (Matsue-shi, Japan), doi: 10.1109/ISLC.2004.1382759.
- Averbeck, R., Jaschke, G., Geelhaar, L., and Riechert, H. (2005). Low threshold InGaAsN/GaAs lasers beyond 1500 nm, *J. Cryst. Growth*, **278**, 224.
- Baba, T., Fukaya, N., and Yonekura, J. (1999). Observation of light propagation in photonic crystal optical waveguides with bends, *Electron. Lett.*, **35**, 654.
- Baier, M., Pelucchi, E., Kapon, E., Varoutsis, S., Gallart, M., Robert-Philip, I., and Abram, I. (2004). Single photon emission from site-controlled pyramidal quantum dots, *Appl. Phys. Lett.*, **84**, 648.
- Baier, M. H., Malko, A., Pelucchi, E., Oberli, D. Y., and Kapon, E. (2006). Quantum-dot exciton dynamics probed by photon-correlation spectroscopy, *Phys. Rev. B*, **73**, 205321.
- Balakrishnan, N., Patanè, A., Makarovskiy, O., Polimeni, A., Capizzi, M., Martelli, F., and Rubini, S. (2011). Laser writing of the electronic activity of N- and H-atoms in GaAs, *Appl. Phys. Lett.*, **99**, 021105.

- Balakrishnan, N., Pettinari, G., Makarovskiy, O., Turyanska, L., Fay, M. W., De Luca, M., Polimeni, A., Capizzi, M., Martelli, F., Rubini, S. and Patanè, A. (2012). Band-gap profiling by laser writing of hydrogen-containing III-N-Vs, *Phys. Rev. B*, **86**, 155307.
- Baldassarri Höger von Högersthal, G., Bissiri, M., Polimeni, A., Capizzi, M., Fischer, M., Reinhardt, M., and Forchel, A. (2001). Hydrogen-induced band gap tuning of (InGa)(AsN)/GaAs single quantum wells, *Appl. Phys. Lett.*, **78**, 3472.
- Baldassarri Höger von Högersthal, G., Polimeni, A., Masia, F., Bissiri, M., Capizzi, M., Gollub, D., Fischer, M., and Forchel, A. (2003). Magnetophotoluminescence studies of (InGa)(AsN)/GaAs heterostructures, *Phys. Rev. B*, **67**, 233304.
- Bank, S. R. (2006). *High-Performance 1.55 μm GaInNAsSb Lasers Grown on GaAs*, PhD thesis, Stanford University.
- Bank, S. R., Bae, H. P., Yuen, H. B., Wistey, M. A., Goddard, L. L., and Harris, J. S. (2006). Room-temperature continuous-wave 1.55 μm GaInNAsSb laser on GaAs, *Electron. Lett.*, **42**, 156.
- Bank, S. R., Wistey, M. A., Goddard, L. L., Yuen, H. B., Lordi, V., and Harris, J. S. (2004). Low-threshold continuous-wave 1.5-μm GaInNAsSb lasers grown on GaAs, *IEEE J. Quantum Electron.*, **40**, 656.
- Bank, S. R., Yuen, H. B., Wistey, M. A., Lordi, V., Bae, H. P., and Harris, J. S. (2005). Effects of growth temperature on the structural and optical properties of 1.55 μm GaInNAsSb quantum wells grown on GaAs, *Appl. Phys. Lett.*, **87**, 21908.
- Barrau, J., Amand, T., Brousseau, M., Simes, R. J., and Goldstein, L. (1992). Induced electrostatic confinement of the electron gas in tensile strained InGaAs/InGaAsP quantum well lasers, *J. Appl. Phys.*, **71**, 5768.
- Barth, M., Nuesse, N., Loechel, B., and Benson, O. (2009). Controlled coupling of a single-diamond nanocrystal to a photonic crystal cavity, *Opt. Lett.*, **34**, 1108.
- Bastard, G. (1990). *Wave Mechanics Applied to Semiconductor Heterostructures* (Editions de Physique, Paris).
- Bayer, M., Gutbrod, T., Reithmaier, J. P., Forchel, A., Reinecke, T. L., Knipp, P. A., Dremin, A. A., and Kulakovskii, V. D. (1998). Optical modes in photonic molecules, *Phys. Rev. Lett.*, **81**, 2582.
- Becke, A. D. (1988). Density-functional exchange-energy approximation with correct asymptotic behavior, *Phys. Rev. A*, **38**, 309.
- Bellaiche, L., Wei, S., and Zunger, A. (1996). Localization and percolation in semiconductor alloys: GaAsN vs GaAsP, *Phys. Rev. B*, **54**, 17568.

- Benfatto, M., and Felici, R. (2001). Resonant atomic scattering factor theory: a multiple scattering approach, *Phys. Rev. B*, **64**, 115410.
- Bergman, K., Stavola, M., Pearton, S. J., and Lopata, J. (1998). Donor-hydrogen complexes in passivated silicon, *Phys. Rev. B*, **37**, 2770.
- Berti, M., Bisognin, G., De Salvador, D., and Napolitani, E. (2008). *Beam Injection Based Nanocharacterization of Advanced Materials* (Research Signpost Trivandrum, Kerala, India).
- Berti, M., Bisognin, G., De Salvador, D., Napolitani, E., Vangelista, S., Polimeni, A., Capizzi, M., Boscherini, F., Ciatto, G., Rubini, S., Martelli, F., and Franciosi, A. (2007). Formation and dissolution of D-N complexes in dilute nitrides, *Phys. Rev. B*, **76**, 205323.
- Bhattacharya, P. (1997). *Semiconductor Optoelectronic Devices* (Prentice Hall, Englewood Cliffs, NJ).
- Biasioli, G., and Kapon, E. (1998). Mechanisms of self-ordering of quantum nanostructures grown on nonplanar surfaces, *Phys. Rev. Lett.*, **81**, 2962.
- Bimberg, D. (1978). Anomaly of the linear and quadratic Zeeman effect of an effective-mass acceptor: C in GaAs, *Phys. Rev. B*, **18**, 1794.
- Bimberg, D., Grundmann, M., and Ledentsov, N. N. (1999). *Quantum Dot Heterostructures* (Wiley, Chichester).
- Bisognin, G., De Salvador, D., Drigo, A. V., Napolitani, E., Sambo, A., Berti, M., Polimeni, A., Felici, M., Capizzi, M., Güngerich, M., Klar, P. J., Bais, G., Jabeen, F., Piccin, M., Rubini, S., Martelli, F. and Franciosi, A. (2006). Hydrogen-nitrogen complexes in dilute nitride alloys: origin of the compressive lattice strain, *Appl. Phys. Lett.*, **89**, 061904.
- Bisognin, G., De Salvador, D., Mattevi, C., Berti, M., Drigo, A. V., Ciatto, G., Grenouillet, L., Duvaut, P., Gilet, P., and Mariette, H. (2004). Determination of lattice parameter and of N lattice location in  $\text{In}_x\text{Ga}_{1-x}\text{N}_y\text{As}_{1-y}/\text{GaAs}$  and  $\text{GaN}_y\text{As}_{1-y}/\text{GaAs}$  epilayers, *J. Appl. Phys.*, **95**, 48.
- Bisognin, G., De Salvador, D., Napolitani, E., Berti, M., Polimeni, A., Capizzi, M., Rubini, S., Martelli, F., and Franciosi, A. (2008). High-resolution x-ray diffraction *in situ* study of very small complexes: the case of hydrogenated dilute nitrides, *J. Appl. Cryst.*, **41**, 366.
- Bisognin, G., De Salvador, D., Napolitani, E., Berti, M., Polimeni, A., Felici, M., Capizzi, M., Bais, G., Jabeen, F., Piccin, M., Rubini, S., Martelli, F., and Franciosi, A. (2007). Thermal evolution of small N-D complexes in deuterated dilute nitrides revealed by *in-situ* high resolution x-ray diffraction, *Phys. Status Solidi A*, **204**, 2766.

- Bissiri, M., Baldassarri Höger von Högersthal, G., Polimeni, A., Capizzi, M., Gollub, D., Fischer, M., Reinhardt, M., and Forchel, A. (2002a). Role of N clusters in  $\text{In}_x\text{Ga}_{1-x}\text{As}_{1-y}\text{N}_y$  band-gap reduction, *Phys. Rev. B*, **66**, 033311.
- Bissiri, M., Baldassarri Höger von Högersthal, G., Polimeni, A., Gaspari, V., Ranalli, F., Capizzi, M., Amore Bonapasta, A., Jiang, F., Stavola, M., Gollub, D., Fischer, M., Reinhardt, M., and Forchel, A. (2002b). Hydrogen-induced passivation of nitrogen in  $\text{GaAs}_{1-y}\text{N}_y$ , *Phys. Rev. B*, **65**, 235210.
- Borenstein, J. T., Corbett, J. W., and Pearton, S. J. (1993). Kinetic-model for hydrogen reactions in boron-doped silicon, *J. Appl. Phys.*, **73**, 2751.
- Bosch, H. S., and Hale, G. M. (1992). Improved formulas for fusion cross-sections and thermal reactivities, *Nucl. Fusion*, **32**, 611.
- Boscherini, F. (2013). X-ray absorption fine structure in the study of semiconductor heterostructures and nanostructures, in *Characterization of Semiconductor Heterostructures and Nanostructures*, 2nd ed. (Elsevier, Amsterdam).
- Bransden, B. H., and Joachain, C. J. (2003). *Physics of Atoms and Molecules* (Pearson Education, Prentice Hall, USA).
- Broderick, C. A., Usman, M., Sweeney, S. J., and O'Reilly, E. P. (2012). Band engineering in dilute nitride and bismide semiconductor lasers, *Semicond. Sci. Technol.*, **27**, 094011.
- Buda, F., Chiarotti, G. L., Car, R., and Parrinello, M. (1989). Proton diffusion in crystalline silicon, *Phys. Rev. Lett.*, **63**, 294.
- Bunker, G. (2010). *Introduction to XAFS: A Practical Guide to X-Ray Absorption Fine Structure Spectroscopy* (Cambridge University Press, UK).
- Burt, M. G. (1999). Fundamentals of envelope function theory for electronic states and photonic modes in nanostructures, *J. Phys. Cond. Matt.*, **11**, R53.
- Buyanova, I. A., and Chen, W. M. (2004). *Physics and Applications of Dilute Nitrides* (Taylor & Francis, New York).
- Buyanova, I. A., Chen, W. M., Izadifard, M., Pearton, S. J., Bihler, C., Brandt, M. S., Hong, Y. G., and Tu, C. W. (2007). Hydrogen passivation of nitrogen in GaNAs and GaNP alloys: how many H atoms are required for each N atom? *Appl. Phys. Lett.*, **90**, 021920.
- Buyanova, I. A., Chen, W. M., Pozina, G., Bergman, J. P., Monemar, B., Xin, H. P., and Tu, C. W. (1999). Mechanism for low-temperature photoluminescence in GaNAs/GaAs structures grown by molecular-beam epitaxy, *Appl. Phys. Lett.*, **75**, 124429.

- Buyanova, I. A., Chen, W. M., Pozina, G., Monemar, B., and Ni, W. X. (1997). Mechanism for thermal quenching of luminescence in SiGe/Si structures grown by molecular beam epitaxy: role of nonradiative defects, *Appl. Phys. Lett.*, **71**, 3676–3678.
- Buyanova, I. A., Chen, W. M., and Tu, C. W. (2003a). Recombination processes in N-containing III-V ternary alloys, *Solid State Electron.*, **47**, 467–475.
- Buyanova, I. A., Ferreira, A. C., Holtz, P. O., Monemar, B., Campman, K., Merz, J. L., and Gossard, A. C. (1996). Effect of hydrogen passivation on Be-doped AlGaAs/GaAs quantum wells, *Appl. Phys. Lett.*, **68**, 1365–1367.
- Buyanova, I. A., Izadifard, M., Chen, W. M., Polimeni, A., Capizzi, M., Xin, H. P., and Tu, C. W. (2003b). Hydrogen-induced improvements in optical quality of GaNAs alloys, *Appl. Phys. Lett.*, **82**, 3662.
- Buyanova, I. A., Izadifard, M., Ivanov, I. G., Birch, J., Chen, W. M., Felici, M., Polimeni, A., Capizzi, M., Hong, Y. G., Xin, H. P., and Tu, C. W. (2004). Direct experimental evidence for unusual effects of hydrogen on the electronic and vibrational properties of  $\text{GaN}_x\text{P}_{1-x}$  alloys: a proof for a general property of dilute nitrides, *Phys. Rev. B*, **70**, 245215.
- Cabib, D., Fabri, E., and Fiorio, G. (1971). The ground state of the exciton in a magnetic field, *Solid State Commun.*, **9**, 1517.
- Cabib, D., Fabri, E., and Fiorio, G. (1972). Ground and first excited states of excitons in a magnetic field, *Il Nuovo Cimento B*, **10**, 185.
- Cai, Y., and Thorpe, M. F. (1992). Length mismatch in random semiconductor alloys. I general theory for quaternaries, *Phys. Rev. B*, **46**, 15872.
- Calic, M., Gallo, P., Felici, M., Atlasov, K. A., Dwir, B., Rudra, A., Biasiol, G., Sorba, L., Tarel, G., Savona, V., and Kapon, E. (2011). Phonon-mediated coupling of InGaAs/GaAs quantum-dot excitons to photonic crystal cavities, *Phys. Rev. Lett.*, **106**, 227402.
- Capizzi, M., Coluzza, C., Emiliani, V., Frankl, P., Frova, A., Sarto, F., Amore Bonapasta, A., Sobiesierski, Z., and Sacks, R. N. (1992). Hydrogen activated radiative states in GaAs/GaAlAs heterostructures and InGaAs/GaAs multiquantum wells, *J. Appl. Phys.*, **72**, 1454–1459.
- Capizzi, M., Emiliani, V., Frova, A., and Sarto, F. (1993). Effect of ion-gun hydrogenation on the photoluminescence of degenerate n-type GaAs:Si, *Phys. Rev. B*, **47**, 4301–4306.
- Caplan, P. J., Poindexter, E. H., Deal, B. E., and Razouk, R. R. (1979). ESR centers, interface states, and oxide fixed charge in thermally oxidized silicon wafers, *J. Appl. Phys.*, **50**, 5847–5854.
- Carrère, H., Marie, X., Lombez, L., and Amand, T. (2006). Optical gain of InGaAsN/InP quantum wells for laser applications, *Appl. Phys. Lett.*, **89**, 181115.

- Carron, R., Fekete, D., Gallo, P., Dwir, B., Rudra, A., Felici, M., Bartova, B., Cantoni, M., and Kapon, E. (2011a). Dilute nitride InGaAsN/GaAs V-groove quantum wires emitting at 1.3  $\mu\text{m}$  wavelength at room temperature, *Appl. Phys. Lett.*, **99**, 101107.
- Carron, R., Gallo, P., Dwir, B., Rudra, A., and Kapon, E. (2011b). Dilute-nitride GaInAsN/GaAs site-controlled pyramidal quantum dots, *Appl. Phys. Lett.*, **99**, 181113.
- Cauchois, Y. (1956). Distribution spectrale observée dans une région d'absorption propre de divers cristaux, in *Compte Rendus de l'Académie des Sciences (Paris)*, **242**, 100–102.
- Chadderton, L. T. (1965). *Radiation Damage in Crystals* (Methuen, London).
- Chakraborty, T., and Pietiläinen, P. (1994). Electron-electron interaction and the persistent current in a quantum ring, *Phys. Rev. B*, **50**, 8460.
- Chen, T. R., Chang, B., Chiu, L. C., Yu, K. L., Margalit, S., and Yariv, A. (1983). Carrier leakage and temperature dependence of InGaAsP lasers, *Appl. Phys. Lett.*, **43**, 217.
- Chen, W. M. (2000). Applications of optically detected magnetic resonance in semiconductor layered structures, *Thin Solid Films*, **364**, 45–52.
- Chen, W. M., Buyanova, I. A., Buyanov, A. V., Lundstrom, T., Bi, W. G., and Tu, C. W. (1996). Intrinsic doping: a new approach for n-type modulation doping in InP-based heterostructures, *Phys. Rev. Lett.*, **77**, 2734–2737.
- Chevallier, J., Dautremont-Smith, W. C., Tu, C. W., and Pearton, S. J. (1985). Donor neutralization in GaAs(Si) by atomic hydrogen, *Appl. Phys. Lett.*, **47**, 108.
- Ciatto, G. (2014). Complexes and clusters in crystalline semiconductors, in *X-Ray Absorption Spectroscopy of Semiconductors* (Springer, Berlin).
- Ciatto, G., Boscherini, F., Amore Bonapasta, A., Filippone, F., Polimeni, A. and Capizzi, M. (2005a). Nitrogen-hydrogen complex in  $\text{GaAs}_x\text{N}_{1-x}$  revealed by x-ray absorption spectroscopy, *Phys. Rev. B*, **71**, 201301(R).
- Ciatto, G., Boscherini, F., Amore Bonapasta, A., Filippone, F., Polimeni, A., Capizzi, M., Berti, M., Bisognin, G., De Salvador, D., Floreano, L., Martelli, F., Rubini, S., and Grenouillet, L. (2009). Local structure of nitrogen-hydrogen complexes in dilute nitrides, *Phys. Rev. B*, **79**, 165205.
- Ciatto, G., d'Acapito, F., Boscherini, F., and Mobilio, S. (2004). Treatment of EXAFS data taken in the fluorescence mode in non-linear conditions, *J. Synchrotron Radiat.*, **11**, 278.
- Ciatto, G., D'Acapito, F., Sanna, S., Fiorentini, V., Polimeni, A., Capizzi, M., Mobilio, S., and Boscherini, F. (2005b). Comparison between

- experimental and theoretical determination of the local structure of the  $\text{GaAs}_{1-y}\text{N}_y$  dilute nitride alloy, *Phys. Rev. B*, **71**, 115210.
- Ciatto, G., Harmand, J.-C., Glas, F., Largeau, L., Le Du, M., Boscherini, F., Malvestuto, M., Floreano, L., Glatzel, P., and Alonso Mori, R. (2007). Anions relative location in the group-V sublattice of  $\text{GaAsSbN}/\text{GaAs}$  epilayers: XAFS measurements and simulations, *Phys. Rev. B*, **75**, 245212.
- Ciatto, G., Renevier, H., Proietti, M. G., Polimeni, A., Capizzi, M., Mobilio, S., and Boscherini, F. (2005c). Effects of hydrogenation on the local structure of  $\text{In}_x\text{Ga}_{1-x}\text{As}_{1-y}\text{N}_y$  quantum wells and  $\text{GaAs}_{1-y}\text{N}_y$  epilayers, *Phys. Rev. B*, **72**, 085322.
- Clerjaud, B., Côte, D., Hahn, W.-S., Lebkiri, A., Ulrici, W., and Wasik, D. (1996). Nitrogen-dihydrogen complex in GaP, *Phys. Rev. Lett.*, **77**, 4930.
- Coccoccioni, M., and de Gironcoli, S. (2005). Linear response approach to the calculation of the effective interaction parameters in the LDA+U method, *Phys. Rev. B*, **71**, 035105.
- Cohen, A. J., Mori-Sánchez, P., and Yang, W. (2008). Insights into current limitations of density functional theory, *Science*, **321**, 792.
- Cohen-Tannoudji, C., Diu, B., and Laloe, F. (1973). *Mécanique Quantique*, Vol. II (Hermann).
- Coleman, J. (1993). Chapter 8, in *Quantum Well Lasers* (Academic Press, New York).
- Cotton, F. A., and Wilkinson, G. (1988). *Advanced Inorganic Chemistry* (John Wiley, New York).
- CPMD (2012). (Copyright IBM Corp 1990-2008, Copyright MPI fuer Festkörperforschung, Stuttgart 1997-2001, <http://www.cpmd.org>).
- D. Marx and J. Hutter (2000). *Ab initio Molecular Dynamics: Theory and Implementation, NIC*, Vol. 1 (John von Neumann Institute for Computing, Jülich), ISBN 3-00-005618, 301.
- D'Acapito, F. (2011). Advanced methods for the analysis of x-ray absorption spectroscopy data applied to semiconductors, *Semic. Science Technol.*, **26**, 064004.
- Dagnelund, D., Tu, C. W., Polimeni, A., Capizzi, M., Chen, W. M., and Buyanova, I. A. (2013). Effect of thermal annealing on defects in post-growth hydrogenated GaNP, *Phys. Status Solidi C*, **10**, 561.
- Dagnelund, D., Vorona, I., Nosenko, G., Wang, X. J., Tu, C. W., Yonezu, H., Polimeni, A., Capizzi, M., Chen, W. M., and Buyanova, I. A. (2012). Effects of hydrogenation on non-radiative defects in GaNP and GaNAs alloys:

- an optically detected magnetic resonance study. *J. Appl. Phys.*, **111**, 023501.
- Dagnelund, D., Wang, X. J., Tu, C. W., Polimeni, A., Capizzi, M., Chen, W. M. and Buyanova, I. A. (2011). Effect of postgrowth hydrogen treatment on defects in GaNP, *Appl. Phys. Lett.*, **98**, 141920.
- Dannecker, T., Jin, Y., Cheng, H., Gorman, C., Buckeridge, J., Uher, C., Fahy, S., Kurdak, C., and Goldman, R. S. (2010). Nitrogen composition dependence of electron effective mass in  $\text{GaAs}_{1-x}\text{N}_x$ , *Phys. Rev. B*, **82**, 125203.
- Davies, G., Lightowlers, E. C., Stavola, M., Bergman, K., and Svensson, B. (1987). The  $3942\text{-cm}^{-1}$  optical band in irradiated silicon, *Phys. Rev. B*, **35**, 2755.
- De Luca, M., Pettinari, G., Ciatto, G., Amidani, L., Filippone, F., Polimeni, A., Fonda, E., Boscherini, F., Amore Bonapasta, A., Giubertoni, D., Knübel, A., Lebedev, V., and Capizzi, M. (2012). Identification of four-hydrogen complexes in In-rich  $\text{In}_x\text{Ga}_{1-x}\text{N}$  ( $x > 0.4$ ) alloys using photoluminescence, x-ray absorption, and density functional theory, *Phys. Rev. B*, **86**, 201202(R).
- Derbyshire, G., Cheung, K. C., Sangsingkeow, P., and Hasnain, S. (1999). A low-profile monolithic multi-element Ge detector for x-ray fluorescence applications, *J. Synchrotron Radiat.*, **6**, 62.
- Diehl, W., Brick, P., Kunert, B., Reinhard, S., Volz, K., and Stoltz, W. (2007). Low threshold 1260 nm (GaIn)(NAs) semiconductor disk laser, *Appl. Phys. Lett.*, **91**, 071103.
- Dieumegard, D., Dubreuil, D., and Amsel, G. (1979). Analysis and depth profiling of deuterium with the  $\text{D}(\text{He}^3, \text{p})\text{He}^4$  reaction by detecting the protons at backward angles, *Nucl. Instrum. Methods*, **166**, 431.
- Dixon, P., Richardson, D., Jones, R., Latham, C. D., Öberg, S., Torres, V. J. B., and Briddon, P. R. (1998). Nitrogen-hydrogen defects in GaP, *Phys. Status Solidi B*, **210**, 321.
- Dovesi, R., Saunders, V. R., Roetti, C., Orlando, R., Zicovich-Wilson, C. M., Pascale, F., Civalleri, B., Doll, K., Harrison, N. M., Bush, I. J., D'Arco, P., and Llunell, M. (2006). *Crystal06 User's Manual*, University of Torino, Torino.
- Du, M.-H., Limpijumnong, S., and Zhang, S. B. (2005). Hydrogen pairs and local vibrational frequencies in H-irradiated  $\text{GaAs}_{1-y}\text{N}_y$ , *Phys. Rev. B*, **72**, 073202.
- Dutta, N. K., and Nelson, R. J. (1981). Temperature dependence of threshold of InGaAsP/InP double heterostructure lasers and Auger recombination, *Appl. Phys. Lett.*, **38**, 407.

- Elias, D. C., Nair, R. R., Mohiuddin, T. M. G., Morozov, S. V., Blake, P., Halsall, M. P., Ferrari, A. C., Boukhvalov, D. W., Katsnelson, M. I., Geim, A. K., and Novoselov, K. (2009). Control of graphene's properties by reversible hydrogenation: evidence for graphane, *Science*, **323**, 610.
- Endicott, J., Patanè, A., Ibáñez, J., Eaves, L., Bissiri, M., Hopkinson, M., Airey, R., and Hill, G. (2003). Magnetotunneling spectroscopy of dilute Ga(AsN) quantum wells, *Phys. Rev. Lett.*, **91**, 126802.
- Englund, D., Fushman, I., and Vuckovic, J. (2005). General recipe for designing photonic crystal cavities, *Opt. Express*, **13**, 5961.
- Englund, D., Shields, B., Rivoire, K., Hatami, F., Vuckovic, J., Park, H. and Lukin, M. D. (2010). Deterministic coupling of a single nitrogen vacancy center to a photonic crystal cavity, *Nano Lett.*, **10**, 3922.
- Erbil, A., Cargill III, G. S., Frahm, R., and Boehme, R. F. (1988). Total-electron-yield current measurements for near-surface extended x-ray-absorption fine structure, *Phys. Rev. B*, **37**, 2450.
- Erol, A. (2008). *Dilute III-V Nitride Semiconductors and Material Systems* (Springer, Berlin).
- Fahy, S., Lindsay, A., Ouerdane, H., and O'Reilly, E. P. (2006). Alloy scattering of n-type carriers in  $\text{GaN}_x\text{As}_{1-x}$ , *Phys. Rev. B*, **74**, 035203.
- Fahy, S., and O'Reilly, E. P. (2003). Intrinsic limits on electron mobility in dilute nitride semiconductors, *Appl. Phys. Lett.*, **83**, 3731.
- Fehse, R., Tomić, S., Adams, A. R., Sweeney, S. J., O'Reilly, E. P., Andreev, A. D., and Riechert, H. (2002). A quantitative study of radiative, Auger, and defect related recombination processes in 1.3- $\mu\text{m}$  GaInNAs-based quantum-well lasers, *IEEE J. Sel. Top. Quantum Electron.*, **8**, 801.
- Felici, M., Atlasov, K. A., Surrente, A., and Kapon, E. (2010). Semianalytical approach to the design of photonic crystal cavities, *Phys. Rev. B*, **82**, 115118.
- Felici, M., Gallo, P., Mohan, A., Dwir, B., Rudra, A., and Kapon, E. (2009). Site-controlled InGaAs quantum dots with tunable emission energy, *Small*, **5**, 938.
- Felici, M., Pettinari, G., Carron, R., Lavenuta, G., Tartaglini, E., Polimeni, A., Fekete, D., Gallo, P., Dwir, B., Rudra, A., Christianen, P. C. M., Maan, J. C., Capizzi, M., and Kapon, E. (2012a). Magneto-optical properties of single site-controlled InGaAsN quantum wires grown on prepatterned GaAs substrates, *Phys. Rev. B*, **85**, 155319.
- Felici, M., Polimeni, A., Miriametro, A., Capizzi, M., Xin, H. P., and Tu, C. W. (2005). Free carrier and/or exciton trapping by nitrogen pairs in dilute  $\text{GaP}_{1-x}\text{N}_x$ , *Phys. Rev. B*, **71**, 045209.

- Felici, M., Polimeni, A., Salviati, G., Lazzarini, L., Armani, N., Masia, F., Capizzi, M., Martelli, F., Lazzarino, M., Bais, G., Piccin, M., Rubini, S. and Franciosi, A. (2006a). In-plane bandgap engineering by modulated hydrogenation of dilute nitride semiconductors, *Adv. Mater.*, **18**, 1993.
- Felici, M., Polimeni, A., Tartaglini, E., Notargiacomo, A., De Luca, M., Carron, R., Fekete, D., Dwir, B., Rudra, A., Capizzi, M., and Kapon, E. (2012b). Reduced temperature sensitivity of the polarization properties of hydrogenated InGaAsN V-groove quantum wires, *Appl. Phys. Lett.*, **101**, 151114.
- Felici, M., Trotta, R., Masia, F., Polimeni, A., Miriametro, A., Capizzi, M., Klar, P. J., and Stoltz, W. (2006b). Compositional disorder in  $\text{GaAs}_{1-x}\text{N}_x:\text{H}$  investigated by photoluminescence, *Phys. Rev. B*, **74**, 085203.
- Filippone, F., Mattioli, G., Alippi, P., and Amore Bonapasta, A. (2011). Clusters and Magnetic Anchoring Points in  $(\text{Ga},\text{Fe})\text{N}$  Condensed Magnetic Semiconductors, *Phys. Rev. Lett.*, **107**, 196401.
- Filippone, F., and Parrinello, M. (2001). Vibrational analysis from linear response theory, *Chem. Phys. Lett.*, **345**, 179.
- Fischer, M., Gollub, D., Reinhardt, M., Kamp, M., and Forchel, A. (2003). GaInNAs for GaAs based lasers for the 1.3 to 1.5  $\mu\text{m}$  range, *J. Cryst. Growth*, **251**, 353.
- Fischer, M., Reinhardt, M., and Forchel, A. (2000). GaInAsN/GaAs laser diodes operating at 1.52  $\mu\text{m}$ , *Electron. Lett.*, **36**, 1208.
- Fisher, M. A., Adams, A. R., O'Reilly, E. P., and Harris, J. J. (1987). Resonant electron scattering due to the central cell of impurities observed in AlGaAs using hydrostatic pressure, *Phys. Rev. Lett.*, **59**, 2341.
- Floreano, L., Naletto, G., Cvetko, D., Gotter, R., Malvezzi, M., Marassi, L., Morgante, A., Santaniello, A., Verdini, A., and Tommasini, F. (1999). Performance of the grating-crystal monochromator of the ALOISA beamline at the Elettra Synchrotron, *Rev. Sci. Instrum.*, **70**, 3855.
- Fowler, B., Martin, K. R., Washer, K., and Stavola, M. (2005). Structure and vibrational properties of N-H<sub>2</sub> complexes in GaAs:N, *Phys. Rev. B*, **72**, 035208.
- Francoeur, S., Klem, J. F., and Mascarenhas, A. (2004). Optical spectroscopy of single impurity centers in semiconductors, *Phys. Rev. Lett.*, **93**, 067403.
- Gallo, P., Felici, M., Dwir, B., Atlasov, K. A., Karlsson, K. F., Rudra, A., Mohan, A., Biasiol, G., Sorba, L., and Kapon, E. (2008). Integration of site-controlled pyramidal quantum dots and photonic crystal membrane cavities, *Appl. Phys. Lett.*, **92**, 263101.

- Geddo, M., Ciabattoni, T., Guizzetti, G., Galli, M., Patrini, M., Polimeni, A., Trotta, R., Capizzi, M., Bais, G., Piccin, M., Rubini, S., Martelli, F. and Franciosi, A. (2007). Photoreflectance and reflectance investigation of deuterium-irradiated GaAsN, *Appl. Phys. Lett.*, **90**, 091907.
- Geddo, M., Patrini, M., Guizzetti, G., Galli, M., Trotta, R., Polimeni, A., Capizzi, M., Martelli, F., and Rubini, S. (2011). Optical study of hydrogen-irradiated GaAsN/GaAs heterostructures, *J. Appl. Phys.*, **109**, 123511.
- Ghiringhelli, G., Tagliaferri, A., Braicovich, L., and Brookes, N. (1998). A simple spherical grating by-pass monochromator dedicated to soft x-ray emission spectroscopy, *Rev. Sci. Instrum.*, **69**, 1610.
- Giannozzi, P., Baroni, S., Bonini, N., Calandra, M., Car, R., Cavazzoni, C., Ceresoli, D., Chiarotti, G. L., Cococcioni, M., Dabo, I., Corso, A. D., Fabris, S., Fratesi, G., de Gironcoli, S., Gebauer, R., Gerstmann, U., Gougaussis, C., Kokalj, A., Lazzeri, M., Martin-Samos, L., Marzari, N., Mauri, F., Mazzarello, R., Paolini, S., Pasquarello, A., Paulatto, L., Sbraccia, C., Scandolo, S., Sclauzero, G., Seitsonen, A. P., Smogunov, A., Umari, P., and Wentzcovitch, R. M. (2009). Quantum Espresso: a modular and open-source software project for quantum simulations of materials, <http://www.quantum-espresso.org>.
- Glatzel, P., and Bergmann, U. (2005). High resolution 1s core hole x-ray spectroscopy in 3d transition metal complexes—electronic and structural information. *Coord. Chem. Rev.*, **249**, 65.
- Goedecker, S., Huetter, J., and Teter, M. (1996). Separable dual-space Gaussian pseudopotentials, *Phys. Rev. B*, **54**, 1703.
- Gonze, X., Stumpf, R., and Scheffler, M. (1991). Analysis of separable potentials, *Phys. Rev. B*, **44**, 8503.
- Granados, D., Garcia, J. M., Ben, T., and Molina, S. I. (2005). Vertical order in stacked layers of self-assembled In(Ga)As quantum rings on GaAs (001), *Appl. Phys. Lett.*, **86**, 071918.
- Greentree, A., Tahan, C., Cole, J., and Hollenberg, L. (2006). Quantum phase transitions of light, *Nat. Phys.*, **2**, 856.
- Groot, F. D., and Kotani, A. (2008). *Core Level Spectroscopy of Solids* (CRC Press, Boca Raton, USA).
- Hague, C. F., Underwood, J. H., Avila, A., Delaunay, R., Ringuenet, H., Marsi, M., and Sacchi, M. (2005). Plane-grating flat-field soft x-ray spectrometer, *Rev. Sci. Instrum.*, **76**, 023110.
- Hahn, W. S., Clerjaud, B., Côte, D., Gendron, F., Porte, C., Ulrici, W., Wasik, D., and Wilkening, W. (1994). *Defects in Semiconductors 17: Nitrogen-Hydrogen Complexes in GaP and GaAs* (Trans Tech, Aedermannsdorf).

- Hai, P. N., Chen, W. M., Buyanova, I. A., Xin, H. P., and Tu, C. W. (2000). Direct determination of electron effective mass in GaNAs/GaAs quantum wells, *Appl. Phys. Lett.*, **77**, 1843.
- Haller, E. E., Hansen, W. L., and Goulding, F. S. (1981). Physics of ultra-pure germanium, *Adv. Phys.*, **30**, 93–138.
- Harris, C., Lindsay, A., and O'Reilly, E. P. (2008). Evolution of N defect states and optical transitions in ordered and disordered  $\text{GaP}_{1-x}\text{N}_x$  alloys, *J. Phys.: Condens. Matter*, **20**, 295211.
- Harrison, J., and Hauser, J. R. (1976). Alloy scattering in ternary III-V compounds, *Phys. Rev. B*, **13**, 5347.
- Hartmann, A., Ducommun, Y., Kapon, E., Hohenester, U., and Molinari, E. (2000). Few-particle effects in semiconductor quantum dots: observation of multicharged excitons, *Phys. Rev. Lett.*, **84**, 5648.
- Hartwigsén, C., Goedecker, S., and Hutter, J. (1998). Relativistic separable dual-space gaussian pseudopotentials from H to Rn, *Phys. Rev. B*, **58**, 3641.
- Henini, M. (2005). *Dilute Nitride Semiconductors* (Elsevier, Amsterdam).
- Hennessy, K., Badolato, A., Winger, M., Gerace, D., Atatüre, M., Gulde, S., Fält, S., Hu, E., and Imamoglu, A. (2007). Quantum nature of a strongly coupled single quantum dot-cavity system, *Nature*, **445**, 896.
- Higashi, T., Sweeney, S. J., Phillips, A. F., Adams, A. R., O'Reilly, E. P., Uchida, T., and Fujii, T. (1999). Experimental analysis of temperature dependence in 1.3- $\mu\text{m}$  AlGaInAs-InP strained MQW lasers, *IEEE J. Sel. Top. Quantum Electron.*, **5**, 413.
- Hjalmarson, H. P., Vogl, P., Wolford, D., and Dow, J. (1980). Theory of substitutional deep traps in covalent semiconductors, *Phys. Rev. Lett.*, **44**, 810.
- Hofmann, M., Wagner, A., Ellmers, C., Schlichenmeier, C., Schafer, S., Höhnsdorf, F., Koch, J., Stolz, W., Koch, S. W., Rühle, W. W., Hader, J., Moloney, J. V., O'Reilly, E. P., Borchert, B., Egorov, A. Y. and Riechert, H. (2001). Gain spectra of (GaIn)(NAs) laser diodes for the 1.3- $\mu\text{m}$ -wavelength regime, *Appl. Phys. Lett.*, **78**, 3009.
- Hofmann, M. R., Gerhardt, N., Wagner, A. M., Ellmers, C., Höhnsdorf, F., Koch, J., Stolz, W., Koch, S. W., Rühle, W. W., Hader, J., Moloney, J. V., O'Reilly, E. P., Borchert, B., Egorov, A. Y., H. Riechert, H. C. S. and Chow, W. W. (2002). Emission dynamics and optical gain of 1.3- $\mu\text{m}$  (GaIn)(NAs)/GaAs lasers. Quantum electronics, *IEEE J. Quantum Electron.*, **38**, 213.
- Hohenberg, P., and Kohn, W. (1964). Inhomogeneous electron gas, *Phys. Rev.*, **136**, B864.

- Holý, V., Pietsch, U., and Baumbach, T. (1999). *High-Resolution X-Ray Scattering from Thin Films and Multilayers* (Springer-Verlag).
- Hudson, A. J., Stevenson, R. M., Bennett, A. J., Young, R. J., Nicoll, C. A., Atkinson, P., Cooper, K., Ritchie, D. A., and Shields, A. J. (2007). Coherence of an entangled exciton-photon state, *Phys. Rev. Lett.*, **99**, 266802.
- Huggenberger, A., Heckelmann, S., Schneider, C., Hoefling, S., Reitzenstein, S., Worschech, L., Kamp, M., and Forchel, A. (2011). Narrow spectral linewidth from single site-controlled In(Ga)As quantum dots with high uniformity, *Appl. Phys. Lett.*, **98**, 131104.
- Hughes, S., Yao, P., Milde, F., Knorr, A., Dalacu, D., Mnaymneh, K., Sazonova, V., Poole, P., Aers, G., Lapointe, J., Cheriton, R., and Williams, R. (2011). Influence of electron-acoustic phonon scattering on off-resonant cavity feeding within a strongly coupled quantum-dot cavity system, *Phys. Rev. B*, **83**, 165313.
- Ignatov, A., Patanè, A., Makarovskiy, O., and Eaves, L. (2006). Terahertz response of hot electrons in dilute nitride Ga(AsN) alloys, *Appl. Phys. Lett.*, **88**, 032107.
- Ikezawa, M., Sakuma, Y., and Masumoto, Y. (2007). Single photon emission from individual nitrogen pairs in GaP, *Jpn. J. Appl. Phys. II*, **46**, L871.
- Intonti, F., Matarazzo, V., Nasir, A., Makarovskiy, O., Campion, R., Patanè, A., Kumar, S., Rastelli, A., Schmidt, O. G., and Gurioli, M. (2011). Nano-sized light emitting diodes by near field laser exposure, *Appl. Phys. Lett.*, **98**, 183102.
- Ivchenko, E. L., Kalevich, V. K., Shiryaev, A. Y., Afanasiev, M. M. and Masumoto, Y. (2010). Optical orientation and spin-dependent recombination in GaAsN alloys under continuous-wave pumping, *J. Phys.: Condens. Matter*, **22**, 465804.
- Jaklevic, J., Kirby, J. A., Klein, M. P., Robertson, A. S., Brown, G. S. and Eisenberger, P. (1977). Fluorescence detection of exafs: sensitivity enhancement for dilute species and thin films, *Sol. State. Commun.*, **23**, 679.
- Janesko, B. G., Henderson, T. M., and Scuseria, G. E. (2009). Screened hybrid density functionals for solid-state chemistry and physics, *Phys. Chem. Chem. Phys.*, **11**, 443.
- Janotti, A., Lyons, J. L., and Van de Walle, C. G. (2012). Hybrid functional calculations of native point defects in InN, *Phys. Status Solidi A*, **209**, 65.

- Janotti, A., Wei, S. H., Zhang, S. B., Kurtz, S., and Van de Walle, C. G. (2003). Interactions between nitrogen, hydrogen, and gallium vacancies in GaAs<sub>1-x</sub>N<sub>x</sub> alloys, *Phys. Rev. B*, **67**, 161201.
- Janotti, A., Zhang, S. B., and Wei, S.-H. (2002a). Hydrogen vibrational modes in GaP:N: the pivotal role of nitrogen in stabilizing the H<sub>2</sub><sup>\*</sup> complex, *Phys. Rev. Lett.*, **88**, 125506.
- Janotti, A., Zhang, S. B., Wei, S.-H., and Van de Walle, C. G. (2002b). Effects of hydrogen on the electronic properties of dilute GaAsN alloys, *Phys. Rev. Lett.*, **89**, 086403.
- Jiang, F., Stavola, M., Capizzi, M., Polimeni, A., Amore Bonapasta, A. and Filippone, F. (2004). Vibrational spectroscopy of hydrogenated GaAs<sub>1-y</sub>N<sub>y</sub>: a structure-sensitive test of an H<sub>2</sub><sup>\*</sup>(N) model, *Phys. Rev. B*, **69**, 041309(R).
- Jimenez-Mier, J., van Ek, J., Ederer, D. L., Callcott, T. A., Jia, J. J., Carlisle, J., Terminello, L., Asfawi, A., and Perera, R. C. (1999). Dynamical behavior of x-ray absorption and scattering at the L edge of titanium compounds: experiment and theory, *Phys. Rev. B*, **59**, 2649.
- Jin, Y. (2010). *Influence of N incorporation on the electronic properties of dilute nitride (In)GaAsN alloys*, Ph.D. thesis, University of Michigan.
- Johnson, N. M., Herring, C., and Chadi, D. J. (1986). Interstitial hydrogen and neutralization of shallow-donor impurities in single-crystal silicon, *Phys. Rev. Lett.*, **56**, 769–772.
- Joly, Y. (2001). X-ray absorption near-edge structure calculations beyond the muffin-tin approximation, *Phys. Rev. B*, **63**, 125120.
- Joly, Y. (2003). Calculating x-ray absorption near-edge structure at very low energy, *J. Synchrotron Radiat.*, **10**, 58.
- Jones, R. E., Yuen, H. B., Liu, T., and Misra, P. (2011). Lattice matchable alloy for solar cells, US Patent Application 20110232730.
- Kane, E. O. (1957). Band structure of indium antimonide, *J. Phys. Chem. Solids*, **1**, 249.
- Kaplyanskii, A. A. (1963). Noncubic centers in cubic crystals and their piezospectroscopic investigation, *Opt. Spectrosc.*, **16**, 329.
- Kapon, E. (1999). *Semiconductor Lasers: I and II* (Academic Press, New York).
- Kapon, E., Hwang, D. M., and Bhat, R. (1989). Stimulated-emission in semiconductor quantum wire heterostructures, *Phys. Rev. Lett.*, **63**, 430.

- Karaiskaj, D., Mascarenhas, A., Klem, J. F., Volz, K., Stolz, W., Adamcyk, M. and Tiedje, T. (2007). Excitons bound to nitrogen pairs in GaAs as seen by photoluminescence of high spectral and spatial resolution, *Phys. Rev. B*, **76**, 12509.
- Kennedy, D. P., and O'Brien, R. R. (1965). Analysis of the impurity atom distribution near the diffusion mask for a planar p-n junction, *IBM J. Res. Dev.*, **9**, 179.
- Kent, P. R. C., Bellaiche, L., and Zunger, A. (2002). Pseudopotential theory of dilute III-V nitrides, *Semicond. Sci. Technol.*, **17**, 851.
- Kent, P. R. C., and Zunger, A. (2001a). Evolution of III-V nitride alloy electronic structure: the localized to delocalized transition, *Phys. Rev. Lett.*, **86**, 2613.
- Kent, P. R. C., and Zunger, A. (2001b). Theory of electronic structure evolution in GaAsN and GaPN alloys, *Phys. Rev. B*, **64**, 115208.
- Khitrova, G., Gibbs, H. M., Kira, M., Koch, S. W., and Scherer, A. (2006). Vacuum Rabi splitting in semiconductors, *Nat. Phys.*, **2**, 81.
- Kilic, C., and Zunger, A. (2002). n-type doping of oxides by hydrogen, *Appl. Phys. Lett.*, **81**, 73.
- Kim, K. S., Yang, G. M., Shim, H. W., Lim, K., Suh, E.-K., and Lee, H. J. (1997). Photoluminescence characterization of biaxial tensile strained GaAs, *J. Appl. Phys.*, **82**, 5103.
- Kim, Y.-S., and Chang, K. J. (2002). Nitrogen-monohydride versus nitrogen-dihydride complexes in GaAs and GaAs<sub>1-x</sub>N<sub>x</sub> alloys, *Phys. Rev. B*, **66**, 073313.
- Kiravittaya, S., Rastelli, A., and Schmidt, O. (2006). Photoluminescence from seeded three-dimensional InAs/GaAs quantum-dot crystals, *Appl. Phys. Lett.*, **88**, 043112.
- Klar, P. J., Grüning, H., Güngerich, M., Heimbrot, W., Koch, J., Torunski, T., Stolz, W., Polimeni, A., and Capizzi, M. (2003). Global changes of the band structure and the crystal lattice of Ga(N,As) due to hydrogenation, *Phys. Rev. B*, **67**, 121206(R).
- Klar, P. J., Grüning, H., Heimbrot, W., Koch, J., Höhnsdorf, F., Stolz, W., Vicente, P. M. A., and Camassel, J. (2000). From N isoelectronic impurities to N-induced bands in the GaN<sub>x</sub>As<sub>1-x</sub> alloy, *Appl. Phys. Lett.*, **76**, 3439.
- Klar, P. J., Grüning, H., Heimbrot, W., Weiser, G., Koch, J., Volz, K., Stolz, W., Koch, S. W., Tomić, S., Choulis, S. A., Hosea, T., O'Reilly, E. P., Hofmann, M., Hader, J., and Moloney, J. V. (2002). Interband transitions of quantum wells and device structures containing Ga(N,As) and (Ga,In)(N, As), *Semicond. Sci. Technol.*, **17**, 830.

- Kleekajai, S., Jiang, F., Colon, K., Stavola, M., Fowler, W. B., Martin, K. R., Polimeni, A., Capizzi, M., Hong, Y. G., Xin, H. P., Tu, C. W., Gais, G., Rubini, S., and Martelli, F. (2008). Vibrational properties of the H–N–H complex in dilute III-N–V alloys: infrared spectroscopy and density functional theory, *Phys. Rev. B*, **77**, 085213.
- Klinger, M. I., Emtsev, V. V., Mashovets, T. V., Ryvkin, S. M., and Vitovskii, N. A. (1981). Ionization mechanism of the defect production in semiconductors. *Radiat. Eff. Defect S.*, **56**, 229–239.
- Kobayashi, N., and Kobayashi, Y. (1991). As and P desorption from III–V semiconductor surface in metalorganic chemical vapor deposition studied by surface photo-absorption, *Jpn. J. Appl. Phys.*, **30**, L1699.
- Kolmogorov, A., Petrovsky, I., and Piscounov, N. (1937). Etude de l' équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique, *Moscow Univ. Bull. Math*, **1**, 1–25.
- Kondow, M., Kitatani, T., Nakatsuka, S., Larson, M. C., Hakahara, K., Yazawa, Y., Okai, M., and Uomi, K. (1997). GaInNAs: a novel material for long-wavelength semiconductor lasers, *IEEE J. Sel. Top. Quantum Electron.*, **3**, 719.
- Kondow, M., Uomi, K., Niwa, A., Kitatani, T., Watahiki, S., and Yazawa, Y. (1996). Gas-source molecular beam epitaxy of  $\text{GaN}_x\text{As}_{1-x}$  using a N radical as the N source, *Jpn. J. Appl. Phys.*, **33**, L1056.
- Koningsberger, D. C., and Prins, R. (1988). *Principles, Applications and Techniques of EXAFS, SEXAFS and XANES* (Wiley, New York).
- Kozlova, N. V., Pettinari, G., Makarovskiy, O., Mori, N., Polimeni, A., Capizzi, M., Zhuang, Q. D., Krier, A., and Patanè, A. (2013). Non-resonant hydrogen dopants in In(AsN): a route to high electron concentrations and mobilities, *Phys. Rev. B*, **87**, 165207.
- Kulik, H. J., Cococcioni, M., Scherlis, D. A., and Marzari, N. (2006). Density functional theory in transition-metal chemistry: a self-consistent Hubbard  $U$  approach, *Phys. Rev. Lett.*, **97**, 103001.
- Kunold, A., Balocchi, A., Zhao, F., Amand, T., Ben Abdallah, N., Harmand, J. and Marie, X. (2011). Giant spin-dependent photo-conductivity in GaAsN dilute nitride semiconductor, *Phys. Rev. B*, **83**, 165202.
- Kuramochi, E., Notomi, M., Mitsugi, S., Shinya, A., Tanabe, T., and Watanabe, T. (2006). Ultrahigh-Q photonic crystal nanocavities realized by the local width modulation of a line defect, *Appl. Phys. Lett.*, **88**, 041112.
- Kurtz, S., Webb, J., Gedvilas, L., Friedman, D., Geisz, J., Olson, J., King, R., Joslin, D., and Karam, N. (2001). Structural changes during annealing of GaInAsN, *Appl. Phys. Lett.*, **78**, 748.

- Landau, L. D., and Lifshitz, E. M. (1986). *Theory of Elasticity* (Pergamon Press).
- Lee, C., Yang, W., and Parr, R. G. (1988). Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density, *Phys. Rev. B*, **37**, 785.
- Lee, P. A., Citrin, P. H., Eisenberger, P., and Kincaid, B. M. (1981). Extended x-ray absorption fine structure: its strengths and limitations as a structural tool, *Rev. Mod. Phys.*, **53**, 769.
- Leifer, K., Pelucchi, E., Watanabe, S., Michelini, F., Dwir, B., and Kapon, E. (2007). Narrow ( $\approx 4$  mev) inhomogeneous broadening and its correlation with confinement potential of pyramidal quantum dot arrays, *Appl. Phys. Lett.*, **91**, 081106.
- Li, L. H., Sallet, V., Patriarche, G., Largeau, L., Bouchoule, S., Travers, L. and Harmand, J. C. (2003). Investigations on GaInNAsSb quinary alloy for  $1.5\text{ }\mu\text{m}$  laser emission on GaAs, *Appl. Phys. Lett.*, **83**, 1298.
- Li, W., Pessa, M., and Likonen, J. (2001). Lattice parameter in GaNAs epilayers on GaAs: deviation from Vegard's law, *Appl. Phys. Lett.*, **78**, 2864.
- Liebich, S., Zimbrich, M., Beyer, A., Lange, C., Franzbach, D. J., Chaterjee, S., Hossain, N., Sweeney, S. J., Volz, K., and Stoltz, W. (2011). Laser operation of Ga(NAsP) lattice-matched to (001) silicon substrate, *Appl. Phys. Lett.*, **99**, 071109.
- Lindsay, A., and O'Reilly, E. P. (1999). Theory of enhanced bandgap non-parabolicity in  $\text{GaN}_x\text{As}_{1-x}$  and related alloys, *Solid State Commun.*, **112**, 443.
- Lindsay, A., and O'Reilly, E. P. (2001). Influence of nitrogen resonant states on the electronic structure of  $\text{GaN}_x\text{As}_{1-x}$ , *Solid State Commun.*, **118**, 313.
- Lindsay, A., and O'Reilly, E. P. (2003). A universal model for trends in  $A_1$ -type defect states in zincblende and diamond semiconductor structures, *Physica B*, **340-342**, 434.
- Lindsay, A., and O'Reilly, E. P. (2004a). A tight-binding-based analysis of the band anti-crossing model in  $\text{GaN}_x\text{As}_{1-x}$ , *Physica E*, **21**, 901.
- Lindsay, A., and O'Reilly, E. P. (2004b). Unification of the band anticrossing and cluster-state models of dilute nitride semiconductor alloys, *Phys. Rev. Lett.*, **93**, 196402.
- Lindsay, A., O'Reilly, E. P., Andreev, A. D., and Ashley, T. (2008). Theory of conduction band structure of  $\text{InN}_x\text{Sb}_{1-x}$  and  $\text{GaN}_x\text{Sb}_{1-x}$  dilute nitride alloys, *Phys. Rev. B*, **77**, 165205.
- Liu, H. Y., Hopkinson, M., Navaretti, P., Gutierrez, M., Ng, J. S., and David, J. P. R. (2003). Improving optical properties of  $1.55\text{ }\mu\text{m}$  GaInNAs/GaAs

- multiple quantum wells with Ga(In)NAs barrier and space layer, *Appl. Phys. Lett.*, **83**, 4951.
- Liu, X., Pistol, M.-E., and Samuelson, L. (1990a). Excitons bound to nitrogen pairs in GaAs, *Phys. Rev. B*, **42**, 7504.
- Liu, X., Pistol, M.-E., Samuelson, L., Schwetlick, S., and Seifert, W. (1990b). Nitrogen pair luminescence in GaAs, *Appl. Phys. Lett.*, **56**, 1451.
- Longa, S. D., Arcovito, A., Girasole, M., Hazemann, J. L., and Benfatto, M. (2001). Quantitative analysis of x-ray absorption near edge structure data by a full multiple scattering procedure: the Fe-CO geometry in photolyzed carbonmonoxy-myoglobin single crystal, *Phys. Rev. Lett.*, **87**, 155501.
- Loridant-Bernard, D., Meziere, S., Constant, M., Dupuy, N., Sombret, S. and Chevalier, J. (1998). Infrared study of light-induced reactivation of neutralized dopants in hydrogenated n-type GaAs doped with silicon, *Appl. Phys. Lett.*, **73**, 644.
- Luo, X. D., Xu, Z. Y., Ge, W. K., Pan, Z., Li, L. H., and Lin, Y. W. (2001). Photoluminescence properties of a  $\text{GaN}_{0.015}\text{As}_{0.985}/\text{GaAs}$  single quantum well under short pulse excitation, *Appl. Phys. Lett.*, **79**, 958.
- Ma, B. S., Su, F. H., Ding, K., Li, G. H., Zhang, Y., Mascarenhas, A., Xin, H. P., and Tu, C. W. (2005). Pressure behavior of the alloy band edge and nitrogen-related centers in  $\text{GaAs}_{0.999}\text{N}_{0.001}$ , *Phys. Rev. B*, **71**, 045213.
- Ma, Y. (1994). X-ray absorption, emission, and resonant inelastic scattering in solids, *Phys. Rev. B*, **49**, 5799.
- Marcet, S., Ouellet-Plamondon, C., Éthier Majcher, G., Saint-Jean, P., André, R., Klem, J., and Francoeur, S. (2010). Charged excitons and biexcitons bound to isoelectronic centers, *Phys. Rev. B*, **82**, 235311.
- Masia, F., Pettinari, G., Polimeni, A., Felici, M., Miriametro, A., Capizzi, M., Lindsay, A., Healy, S. B., O'Reilly, E. P., Cristofoli, A., Bais, G., Piccin, M., Rubini, S., Martelli, F., Franciosi, A., Klar, P. J., Volz, K. and Stolz, W. (2006). Interaction between conduction band edge and nitrogen states probed by carrier effective-mass measurements in  $\text{GaAs}_{1-x}\text{N}_x$ , *Phys. Rev. B*, **73**, 073201.
- Masia, F., Polimeni, A., Baldassarri Höger von Högersthal, G., M. Bissiri, Capizzi, M., Klar, P. J., and Stolz, W. (2003). Early manifestation of localization effects in diluted  $\text{Ga}(\text{AsN})$ , *Appl. Phys. Lett.*, **82**, 4474.
- Massies, J., and Grandjean, N. (1993). Surfactant effect on the surface diffusion length in epitaxial growth, *Phys. Rev. B*, **48**, 8502.
- Matsubara, M., Godet, J., and Pizzagalli, L. (2010). Theoretical study of hydrogen stability and aggregation in dislocation cores in silicon, *Phys. Rev. B*, **82**, 024107.

- Mattila, T., Wei, S. H., and Zunger, A. (1999). Localization and anticrossing of electron levels in GaAs<sub>1-x</sub>N<sub>x</sub> alloys, *Phys. Rev. B*, **60**, R11245.
- Mattioli, G., Alippi, P., Filippone, F., Caminiti, R., and Amore Bonapasta, A. (2010). Deep versus shallow behavior of intrinsic defects in rutile and anatase TiO<sub>2</sub> polymorphs, *J. Phys. Chem. C*, **114**, 21694.
- McCluskey, M. D. (2000). Local vibrational modes of impurities in semiconductors, *J. Appl. Phys.*, **87**, 3593.
- McCluskey, M. D., and Haller, E. E. (1999). Hydrogen in III-V and II-VI semiconductors, in *Hydrogen in Semiconductors II* (Academic Press).
- Merz, J., Faulkner, R., and Dean, P. (1969). Excitonic molecule bound to the isoelectronic nitrogen trap in GaP, *Phys. Rev.*, **188**, 1228.
- Mohan, A., Gallo, P., Felici, M., Dwir, B., Rudra, A., Faist, J., and Kapon, E. (2010). Record-low inhomogeneous broadening of site-controlled quantum dots for nanophotonics, *Small*, **6**, 1268.
- Moller, W., and Besembacher, F. (1980). A note on the He+D nuclear-reaction cross section, *Nucl. Instrum. Methods*, **168**, 111.
- Morrow, R. A. (1989). Modeling the diffusion of hydrogen in GaAs, *J. Appl. Phys.*, **66**, 2973.
- Naber, J. A., and James, H. M. (1961). *Bull. Am. Phys. Soc.*, **6**, 303.
- Nagamune, Y., Arakawa, Y., Tsukamoto, S., Nishioka, M., Sasaki, S., and Miura, N. (1992). Photoluminescence spectra and anisotropic energy shift of GaAs quantum wires in high magnetic-fields, *Phys. Rev. Lett.*, **69**, 2963.
- Newville, M., Ravel, B., Haskel, D., Rehr, J. J., Stern, E. A., and Yacoby, Y. (1995). Analysis of multiple-scattering XAFS data using theoretical standards, *Physica B*, **208&209**, 154.
- Noda, S. (2006). Seeking the ultimate nanolaser, *Science*, **314**, 260.
- Nordgren, J., Bray, G., Gamn, S., Nyholm, R., Rubensson, J., and Wassdahl, N. (1989). Soft x-ray emission spectroscopy using monochromatized synchrotron radiation, *Rev. Sci. Instrum.*, **60**, 1690.
- Northrup, J. E. (2002). Theory of intrinsic and H-passivated screw dislocations in GaN, *Phys. Rev. B*, **66**, 045204.
- Oestreich, M., and Rühle, W. W. (1995). Temperature dependence of the electron landé g factor in GaAs, *Phys. Rev. Lett.*, **74**, 2315.
- O'Reilly, E. P. (1989). Valence band engineering in strained-layer structures, *Semicond. Sci. Technol.*, **4**, 121.
- O'Reilly, E. P., and Lindsay, A. (2010). Tight-binding approach to calculation of localised perturbations in semiconductors, *J. Phys.: Conf. Ser.*, **242**, 012002.

- O'Reilly, E. P., Lindsay, A., and Fahy, S. (2004). Theory of the electronic structure of dilute nitride alloys: beyond the band-anti-crossing model, *J. Phys.: Condens. Matter*, **16**, S3257.
- O'Reilly, E. P., Lindsay, A., Klar, P. J., Polimeni, A., and Capizzi, M. (2009). Trends in the electronic structure of dilute nitride alloys, *Semicond. Sci. Technol.*, **24**, 033001.
- O'Reilly, E. P., Lindsay, A., Tomić, S., and Kamal-Saadi, M. (2002). Tight-binding and k-p models for the electronic structure of Ga(In)NAs and related alloys, *Semicond. Sci. Technol.*, **17**, 870.
- Oyanagi, H., Martini, M., and Saito, M. (1998). Nineteen-element high-purity Ge solid-state detector array for fluorescence x-ray absorption fine structure studies, *Nucl. Instrum. Meth. A*, **403**, 58.
- Pan, J. W., and Chyi, J. I. (1996). Theoretical study of the temperature dependence of 1.3-μm AlGaInAs-InP multiple-quantum-well lasers, *IEEE J. Quantum Electron.*, **32**, 2133.
- Pankove, J. I., Carlson, D. E., Berkeyheiser, J. E., and Wance, R. O. (1983). Neutralization of shallow acceptor levels in silicon by atomic-hydrogen, *Phys. Rev. Lett.*, **51**, 2224–2225.
- Pankove, J. I., and Johnson, N. M. (1991). *Hydrogen in Semiconductors, in Semiconductors and Semimetals*, Vol. 34 (Academic Press, Boston).
- Pauling, L. (1967). *The Nature of the Chemical Bond* (Cornell University Press, Ithaca).
- Pavesi, L., and Giannozzi, P. (1991). H passivation of Si impurities in GaAs, *Phys. Rev. B*, **43**, 2446.
- Pavesi, L., and Giannozzi, P. (1992). Atomic and molecular hydrogen in gallium arsenide: a theoretical study, *Phys. Rev. B*, **46**, 4621.
- Pavesi, L., and Guzzi, M. (1994). Photoluminescence of  $\text{Al}_x\text{Ga}_{1-x}\text{As}$  alloys, *J. Appl. Phys.*, **75**, 4779.
- Pavesi, M., Manfredi, M., Salviati, G., Armani, N., Rossi, F., Meneghesso, G., Levada, S., Zanoni, E., Du, S., and Eliashevich, I. (2004). Optical evidence of an electrothermal degradation of InGaN-based light-emitting diodes during electrical stress, *Appl. Phys. Lett.*, **84**, 3403.
- Pearson, S. J. (1994). Hydrogen in crystalline semiconductors: part II; III-V compounds, *Int. J. Mod. Phys. B*, **8**, 1247.
- Pearson, S. J., Corbett, J. W., and Stavola, M. (1992). *Hydrogen in Crystalline Semiconductors* (Springer-Verlag, Berlin).
- Pearson, S. J., Dautremont-Smith, W. C., Chevallier, J., Tu, C. W. and Cummings, K. D. (1986). Hydrogenation of shallow-donor levels in GaAs, *J. Appl. Phys.*, **59**, 2821–2827.

- People, R., and Bean, J. C. (1985). Calculation of critical layer thickness versus lattice mismatch for  $\text{Ge}_x\text{Si}_{1-x}/\text{Si}$  strained-layer heterostructures, *Appl. Phys. Lett.*, **47**, 322.
- Perkins, J. D., Mascarenhas, A., Zhang, Y., Geisz, J. F., Friedman, D. J., Olson, J. M., and Kurtz, S. R. (1999). Nitrogen-activated transitions, level repulsion, and band gap reduction in  $\text{GaAs}_{1-x}\text{N}_x$  with  $x < 0.03$ , *Phys. Rev. Lett.*, **82**, 3312.
- Peter, E., Senellart, P., Martrou, D., Lemaître, A., Hours, J., Gérard, J., and Bloch, J. (2005). Exciton-photon strong-coupling regime for a single quantum dot embedded in a microcavity, *Phys. Rev. Lett.*, **95**, 067401.
- Pettinari, G., Masia, F., Capizzi, M., Polimeni, A., Losurdo, M., Bruno, G., Kim, T. H., Choi, S., Brown, A., Lebedev, V., Cimalla, V., and Ambacher, O. (2008). Experimental evidence of different hydrogen donors in n-type InN, *Phys. Rev. B*, **77**, 125207.
- Pettinari, G., Masia, F., Polimeni, A., Felici, M., Frova, A., Capizzi, M., Lindsay, A., O'Reilly, E. P., Klar, P. J., Stolz, W., Bais, G., Piccin, M., Rubini, S., Martelli, F., and Franciosi, A. (2006). Influence of nitrogen-cluster states on the gyromagnetic factor of electrons in  $\text{GaAs}_{1-x}\text{N}_x$ , *Phys. Rev. B*, **74**, 245202.
- Pettinari, G., Patanè, A., Polimeni, A., Capizzi, M., Lu, X., and Tiedje, T. (2012a). Effects of hydrogen on the electronic properties of Ga(AsBi) alloys, *Appl. Phys. Lett.*, **101**, 222103.
- Pettinari, G., Polimeni, A., and Capizzi, M. (2012b). *Photoluminescence: a tool for investigating optical, electronic, and structural properties of semiconductors, Chapter 5 in Semiconductor Research* (Spring-Verlag, Berlin, Heidelberg).
- Pettinari, G., Polimeni, A., Masia, F., Trotta, R., Felici, M., Capizzi, M., Niebling, T., Stolz, W., and Klar, P. J. (2007). Electron mass in dilute nitrides and its anomalous dependence on hydrostatic pressure, *Phys. Rev. Lett.*, **98**, 146402.
- Phillips, A. F., Sweeney, S. J., Adams, A. R., and Thijs, P. J. A. (1999). The temperature dependence of 1.3- and 1.5- $\mu\text{m}$  compressively strained InGaAs(P) MQW semiconductor lasers, *IEEE J. Sel. Top. Quantum Electron.*, **5**, 401.
- Phillips, J. C. (1973). *Bonds and Bands in Semiconductors* (Academic Press, New York).
- Polimeni, A. (2007). private communication.
- Polimeni, A., Baldassarri Höger von Högersthal, G., Bissiri, M., Capizzi, M., Fischer, M., Reinhardt, M., and Forchel, A. (2001a). Effect of hydrogen on the electronic properties of  $\text{In}_x\text{Ga}_{1-x}\text{As}_{1-y}\text{N}_y/\text{GaAs}$  quantum wells, *Phys. Rev. B*, **63**, 201304R.

- Polimeni, A., Baldassarri Höger von Högersthal, G., Bissiri, M., Capizzi, M., Frova, A., Fischer, M., Reinhardt, M., and Forchel, A. (2002). Role of hydrogen in III-N-V compound semiconductors, *Semicond. Sci. Technol.*, **17**, 797.
- Polimeni, A., Baldassarri Höger von Högersthal, G., Masia, F., Frova, A., Capizzi, M., Sanna, S., Fiorentini, V., Klar, P. J., and Stoltz, W. (2004). Tunable variation of the electron effective mass and exciton radius in hydrogenated  $\text{GaAs}_{1-x}\text{N}_x$ , *Phys. Rev. B*, **69**, 041201(R).
- Polimeni, A., Bissiri, M., Felici, M., Capizzi, M., Buyanova, I. A., Chen, W. M., Xin, H. P., and Tu, C. W. (2003a). Nitrogen passivation induced by atomic hydrogen: the  $\text{GaP}_{1-y}\text{N}_y$  case, *Phys. Rev. B*, **67**, 201303.
- Polimeni, A., Capizzi, M., Geddo, M., Fischer, M., Reinhardt, M., and Forchel, A. (2001b). Effect of nitrogen on the temperature dependence of the energy gap in  $\text{In}_x\text{Ga}_{1-x}\text{As}_{1-y}\text{N}_y/\text{GaAs}$  single quantum wells, *Phys. Rev. B*, **63**, 195320.
- Polimeni, A., Ciatto, G., Ortega, L., Jiang, F., Boscherini, F., Filippone, F., Amore Bonapasta, A., Stavola, M., and Capizzi, M. (2003b). Lattice relaxation by atomic hydrogen irradiation of III-N-V semiconductor alloys, *Phys. Rev. B*, **68**, 085204.
- Polimeni, A., Marangio, D., Capizzi, M., Frova, A., and Martelli, F. (1994). Giant photoluminescence enhancement in deuterated highly strained InAs/GaAs quantum wells, *Appl. Phys. Lett.*, **65**, 1254.
- Polimeni, A., Masia, F., Pettinari, G., Trotta, R., Felici, M., Capizzi, M., Lindsay, A., O'Reilly, E. P., Niebling, T., Stoltz, W., and Klar, P. J. (2008a). Role of strain and properties of N clusters at the onset of the alloy limit in  $\text{GaAs}_{1-x}\text{N}_x$ , *Phys. Rev. B*, **77**, 155213.
- Polimeni, A., Pettinari, G., Trotta, R., Masia, F., Felici, M., Capizzi, M., Lindsay, A., O'Reilly, E. P., Niebling, T., Stoltz, W., Klar, P. J., Martelli, F., and Rubini, S. (2008b). Photoluminescence under magnetic field and hydrostatic pressure for probing the electronic properties of GaAsN, *Phys. Status Solidi A*, **205**, 107.
- Pollak, F., and Cardona, M. (1968). Piezo-electroreflectance in Ge, GaAs, and Si, *Phys. Rev.*, **172**, 816.
- Proietti, M. G., Renevier, H., Hodeau, J. L., Garcia, J., Berar, J. F. and Wolfers, P. (1999). Diffraction-anomalous-fine-structure spectroscopy applied to the study of III-V strained semiconductors, *Phys. Rev. B*, **59**, 5479.
- Prokofyeva, T., Sauncy, T., Seon, M., Holtz, M., Qiu, Y., Nikishin, S. and Temkin, H. (1998). Raman studies of nitrogen incorporation in  $\text{GaAs}_{1-x}\text{N}_x$ , *Appl. Phys. Lett.*, **73**, 1409.

- Putrino, A., Sebastiani, D., and Parrinello, M. (2000). Generalized variational density functional perturbation theory, *J. Chem. Phys.*, **113**, 7102.
- Raisanen, J., Keinonen, J., and Karttunen, V. (1988). Redistribution of implanted H in annealings of n-type GaAs, *J. Appl. Phys.*, **64**, 2334.
- Rammohan, K., Tang, Y., Rich, D. H., Goldman, R. S., Wieder, H. H. and Kavanagh, K. L. (1995). Relaxation-induced polarized luminescence from  $\text{In}_x\text{Ga}_{1-x}\text{As}$  films grown on GaAs(001), *Phys. Rev. B*, **51**, 5033.
- Reithmaier, J. P., Sek, G., Löffler, A., Hofmann, C., Kuhn, S., Reitzenstein, S., Keldysh, L. V., Kulakovskii, V. D., Reinecke, T. L. and Forchel, A. (2004). Strong coupling in a single quantum dot-semiconductor microcavity system, *Nature*, **432**, 197.
- Renevier, H., Grenier, S., Arnaud, S., Berar, J. F., Caillot, B., Hodeau, J. L., Letoublon, A., Proietti, M. G., and Ravel, B. (2003). Diffraction anomalous fine-structure spectroscopy at beamline BM2 at the european synchrotron radiation facility, *J. Synchrotron Rad.*, **10**, 435.
- Riechert, H., Ramakrishnan, A., and Steinle, G. (2002). Development of InGaAsN-based 1.3  $\mu\text{m}$  VCSELs, *Semicond. Sci. Technol.*, **17**, 892.
- Rossi, J., Wolfe, C., and Dimmock, J. (1970). Acceptor luminescence in high-purity n-type GaAs, *Phys. Rev. Lett.*, **25**, 1614.
- Roth, L. M., Lax, B., and Zwerdling, S. (1959). Theory of optical magneto-absorption effects in semiconductors, *Phys. Rev.*, **114**, 90.
- Rühle, W., and Göbel, E. (1976). New aspects of the magnetoluminescence of a band to acceptor transition in GaAs, *Phys. Status Solidi B*, **78**, 311.
- Rutherford, E. (1911). The scattering of  $\alpha$  and  $\beta$  particles by matter and the structure of the atom, *Philos. Mag.*, **6**, 21.
- Sah, C. T., Sun, J. Y., and Tzou, J. J. (1983). Deactivation of the boron acceptor in silicon by hydrogen, *Appl. Phys. Lett.*, **43**, 204–206.
- Sakurai, J. J. (1967). *Advanced Quantum Mechanics* (Benjamin Cummings).
- Salter, C. L., Stevenson, R. M., Farrer, I., Nicoll, C. A., Ritchie, D. A. and Shields, A. J. (2010). An entangled-light-emitting diode, *Nature*, **465**, 594.
- Sankey, O. F., Dow, J. D., and Hess, K. (1982). Theory of resonant scattering in semiconductors due to impurity central-cell potentials, *Appl. Phys. Lett.*, **41**, 664.
- Sanna, S., and Fiorentini, V. (2004). Lattice constant, effective mass, and gap recovery in hydrogenated  $\text{GaAs}_{1-x}\text{N}_x$ , *Phys. Rev. B*, **69**, 125208.
- Santos, P. V., Johnson, N. M., and Street, R. A. (1991). Light-enhanced hydrogen motion in a-Si:H, *Phys. Rev. Lett.*, **67**, 2686.

- Sayers, D. E., Stern, E. A., and Lytle, F. W. (1971). New technique for investigating noncrystalline structures: Fourier analysis of the extended x-ray-absorption fine structure, *Phys. Rev. Lett.*, **27**, 1204.
- Schaich, W. L. (1973). Comment on the theory of extended x-ray-absorption fine structure, *Phys. Rev. B*, **8**, 4028.
- Schnohr, C., and Ridgway, M. (2014). *X-Ray Absorption Spectroscopy of Semiconductors* (Springer, Berlin).
- Seki, S., Yamanaka, T., Lui, W., Yoshikuni, Y., and Yokoyama, K. (1994). Theoretical-analysis of pure effects of strain and quantum confinement on differential gain in InGaAsP/InP strained-layer quantum-well lasers, *IEEE J. Quantum Electron.*, **30**, 500.
- Shafi, M., Mari, R. H., Khatab, A., Henini, M., Polimeni, A., Capizzi, M. and Hopkinson, M. (2011). Deep levels in H-irradiated GaAs<sub>1-x</sub>N<sub>x</sub> ( $x < 0.01$ ) grown by molecular beam epitaxy, *J. Appl. Phys.*, **110**, 124508.
- Shan, W., Walukiewicz, W., Ager III, J. W., Haller, E. E., Geisz, J. F., Friedman, D. J., Olson, J. M., and Kurtz, S. R. (1999a). Band anticrossing in GaInNAs alloys, *Phys. Rev. Lett.*, **82**, 1221.
- Shan, W., Walukiewicz, W., Ager III, J. W., Haller, E. E., Geisz, J. F., Friedman, D. J., Olson, J. M., and Kurtz, S. R. (1999b). Effect of nitrogen on the band structure of GaInNAs alloys, *J. Appl. Phys.*, **86**, 2349.
- Shantharama, L. G., Adams, A. R., Ahmad, C. N., and Nicholas, R. J. (1984). The  $k \cdot p$  interaction in InP and GaAs from the band-gap dependence of the effective mass, *J. Phys. C*, **17**, 4429.
- Silvestre, S., Bernard-Loridan, D., Constant, E., Constant, M., and Chevallier, J. (2000). Electron-beam-induced reactivation of Si dopants in hydrogenated GaAs: a minority carrier generation effect or an energetic electron excitation effect? *Appl. Phys. Lett.*, **77**, 3206.
- Skiba-Szymanska, J., Jamil, A., Farrer, I., Ward, M. B., Nicoll, C. A., Ellis, D. J. P., Griffiths, J. P., Anderson, D., Jones, G. A. C., Ritchie, D. A. and Shields, A. J. (2011). Narrow emission linewidths of positioned InAs quantum dots grown on pre-patterned GaAs(100) substrates, *Nanotechnology*, **22**, 065302.
- Skierbiszewski, C. (2002). Experimental studies of the conduction band structure of GaInNAs alloys, *Semicond. Sci. Technol.*, **17**, 803.
- Song, B. S., Noda, S., Asano, T., and Akahane, Y. (2005). Ultra-high-Q photonic double-heterostructure nanocavity, *Nat. Mater.*, **4**, 207.
- Spasov, S., Allison, G., Patanè, A., Eaves, L., Tretyakov, M. Y., Ignatov, A., Hopkinson, M., and Hill, G. (2008). High field electron dynamics in dilute nitride Ga(AsN), *Appl. Phys. Lett.*, **93**, 022111.

- Srinivasan, K., and Painter, O. (2003). Fourier space design of high-Q cavities in standard and compressed hexagonal lattice photonic crystals, *Opt. Express*, **11**, 579.
- Stavola, M. (1998). Vibrational spectroscopy of light element impurities in semiconductors, in *Semiconductors and Semimetals 51B* (Academic Press, Boston).
- Stragier, H., Cross, J. O., Rehr, J. J., Sorensen, L. B., Bouldin, C. E. and Woicik, J. C. (1992). Diffraction anomalous fine structure: a new x-ray structural technique, *Phys. Rev. Lett.*, **69**, 3064.
- Strocov, V. N., Nilsson, P. O., Schmitt, T., Augustsson, A., Gridneva, L., Debowska-Nilsson, D., Claessen, R., Egorov, A. Y., Ustinov, V. M. and Alferov, Z. I. (2004). Nitrogen local electronic structure in Ga(In)AsN alloys by soft-x-ray absorption and emission: implications for optical properties, *Phys. Rev. B*, **69**, 035206.
- Suemune, I., Uesugi, K., and Walukiewicz, W. (2000). Role of nitrogen in the reduced temperature dependence of band-gap energy in GaNAs, *Appl. Phys. Lett.*, **77**, 3021.
- Surrente, A., Gallo, P., Felici, M., Dwir, B., Rudra, A., and Kapon, E. (2009). Dense arrays of ordered pyramidal quantum dots with narrow linewidth photoluminescence spectra, *Nanotechnology*, **20**, 415205.
- Tansu, N., Kirsch, N. J., and Mawst, L. J. (2002). Low-threshold-current-density 1300-nm dilute-nitride quantum well lasers, *Physica E*, **81**, 2523.
- Tansu, N., and Mawst, L. J. (2002). Temperature sensitivity of 1300-nm InGaAsN quantum-well lasers, *IEEE Photon. Technol. Lett.*, **14**, 1052.
- Teo, B. K., and Joy, D. C. (1981). *EXAFS Spectroscopy: Techniques and Applications* (Plenum, New York).
- Tesmer, J. R., and Nastasi, M. (1995). *Handbook of Modern Ion Beam Materials Analysis* (Materials Research Society, Pittsburgh).
- Teukam, Z., Chevallier, J., Saguy, C., Kalish, R., Ballutaud, D., Barbé, M., Jomard, F., Tromson-Carli, A., Cytermann, C., Butler, J. E., Bernard, M., Baron, C., and Deneuville, A. (2003). Shallow donors with high n-type electrical conductivity in homoepitaxial deuterated boron-doped diamond layers, *Nat. Mater.*, **2**, 482.
- Thinh, N. Q., Buyanova, I. A., Hai, P. N., Chen, W. M., Xin, H. P., and Tu, C. W. (2001). Signature of an intrinsic point defect in  $\text{GaN}_x\text{As}_{1-x}$ , *Phys. Rev. B*, **63**, 033203.
- Thomas, D. G., and Hopfield, J. J. (1966). Isoelectronic traps due to nitrogen in gallium phosphide, *Phys. Rev.*, **150**, 680.

- Tisch, U., Finkman, E., and Salzman, J. (2002). The anomalous bandgap bowing in GaAsN, *Appl. Phys. Lett.*, **81**, 463.
- Togan, E., Chu, Y., Trifonov, A. S., Jiang, L., Maze, J., Childress, L., Dutt, M. V. G., Sørensen, A. S., Hemmer, P. R., Zibrov, A. S., and Lukin, M. D. (2010). Quantum entanglement between an optical photon and a solid-state spin qubit, *Nature*, **466**, 730.
- Tomić, S., and O'Reilly, E. P. (2002). Gain characteristics of ideal dilute nitride quantum well lasers, *Physica E*, **12**, 1102.
- Tomić, S., and O'Reilly, E. P. (2003). Optimization of material parameters in 1.3- $\mu\text{m}$  InGaAsN-GaAs lasers, *IEEE Photon. Technol. Lett.*, **15**, 6.
- Tomić, S., O'Reilly, E. P., Fehse, R., Sweeney, S. J., Adams, A. R., Andreev, A. D., Choulis, S. A., Hosea, T. J. C., and Riechert, H. (2003). Theoretical and experimental analysis of 1.3- $\mu\text{m}$  InGaAsN/GaAs lasers, *IEEE J. Sel. Top. Quantum Electron.*, **9**, 1228.
- Tomić, S., O'Reilly, E. P., Klar, P. J., Grüning, H., Heimbrodt, W., Chen, W. M., and Buyanova, I. A. (2004). Influence of conduction-band nonparabolicity on electron confinement and effective mass in  $\text{GaN}_x\text{As}_{1-x}/\text{GaAs}$  quantum wells, *Phys. Rev. B*, **69**, 245305.
- Tomljenovic-Hanic, S., Greentree, A. D., De Sterke, C. M., and Prawer, S. (2009). Flexible design of ultrahigh-Q microcavities in diamond-based photonic crystal slabs, *Opt. Express*, **17**, 6465.
- Tormen, M., De Salvador, D., Natali, M., Drigo, A., Romanato, F., Rossetto, G., Boscherini, F., and Mobilio, S. (1999). Bond length variation in  $\text{In}_{0.25}\text{Ga}_{0.75}\text{As}/\text{InP}$  epitaxial layers thicker than the critical thickness, *J. Appl. Phys.*, **86**, 2533.
- Trotta, R., Atkinson, P., Plumhof, J. D., Zallo, E., Rezaev, R. O., Kumar, S., Baunack, S., Schroter, J. R., Rastelli, A., and Schmidt, O. G. (2012a). Nanomembrane quantum-light-emitting diodes integrated onto piezoelectric actuators, *Adv. Mater.*, **24**, 2668.
- Trotta, R., Cavigli, L., Felisari, L., Polimeni, A., Vinattieri, A., Gurioli, M., Capizzi, M., Martelli, F., Rubini, S., Mariucci, L., Francardi, M. and Gerardino, A. (2010). Quantum confinement effects in hydrogen-intercalated  $\text{Ga}_{1-x}\text{As}_x\text{N}_x\text{-GaAs}_{1-x}\text{N}_x$ : H planar heterostructures investigated by photoluminescence spectroscopy, *Phys. Rev. B*, **81**, 235327.
- Trotta, R., Giubertoni, D., Polimeni, A., Bersani, M., Capizzi, M., Martelli, F., Rubini, S., Bisognin, G., and Berti, M. (2009a). Hydrogen diffusion in  $\text{GaAs}_{1-x}\text{N}_x$ , *Phys. Rev. B*, **80**, 195206.

- Trotta, R., Polimeni, A., and Capizzi, M. (2012b). Hydrogen incorporation in III-N-V semiconductors: from macroscopic to nanometre control of the material physical properties, *Adv. Funct. Mater.*, **22**, 1782.
- Trotta, R., Polimeni, A., Capizzi, M., Giubertoni, D., Bersani, M., Bisognin, G., Berti, M., Rubini, S., Martelli, F., Mariucci, L., Francardi, M. and Gerardino, A. (2008). Effect of hydrogen incorporation temperature in in-plane engineered GaAsN/GaAsN:H heterostructure, *Appl. Phys. Lett.*, **92**, 221901.
- Trotta, R., Polimeni, A., Capizzi, M., Martelli, F., Rubini, S., Francardi, M., Gerardino, A., and Mariucci, L. (2009b). Light polarization control in strain-engineered GaAsN/GaAsN:H heterostructures, *Appl. Phys. Lett.*, **94**, 261905.
- Trotta, R., Polimeni, A., Martelli, F., Pettinari, G., Capizzi, M., Felisari, L., Rubini, S., Francardi, M., Gerardino, A., Christianen, P. C. M., and Maan, J. C. (2011). Fabrication of site-controlled quantum dots by spatially selective incorporation of hydrogen in Ga(AsN)/GaAs heterostructures, *Adv. Mater.*, **23**, 2706.
- Trotta, R., Zallo, E., Ortix, C., Atkinson, P., Plumhof, J. D., den Brink, J. V., Rastelli, A., and Schmidt, O. G. (2012c). Universal recovery of the energy-level degeneracy of bright excitons in InGaAs quantum dots without a structure symmetry, *Phys. Rev. Lett.*, **109**, 147401.
- Troullier, N., and Martins, J. (1991). Efficient pseudopotentials for plane-wave calculations, *Phys. Rev. B*, **43**, 1993.
- Ugliengo, P. (2006). *MOLDRAW Program to display and manipulate molecular and crystal structures*, University of Torino, Torino, available at <http://www.moldraw.unito.it>.
- Ugliengo, P., Viterbo, D., and Chiari, G. (1993). MOLDRAW: molecular graphics on a personal computer, *Z. Kristallogr.*, **207**, 9.
- Ungaro, G., Roux, G. L., Teissier, R., and Harmand, J. C. (1999). GaAsSbN: a new low-bandgap material for GaAs substrates, *Electron. Lett.*, **35**, 1246.
- Van de Walle, C. G., Denteneer, P. J. H., Bar-Yam, Y., and Pantelides, S. T. (1989). Theory of hydrogen diffusion and reactions in crystalline silicon, *Phys. Rev. B*, **39**, 10791.
- Van de Walle, C. G., Limpijumnong, S., and Neugebauer, J. (2001). First-principles studies of beryllium doping of GaN, *Phys. Rev. B*, **63**, 245205.
- Van de Walle, C. G., Lyons, J. L., and Janotti, A. (2010). Controlling the conductivity of InN, *Phys. Status Solidi A*, **207**, 1024.

- Van de Walle, C. G., and Neugebauer, J. (2003). Universal alignment of hydrogen levels in semiconductors, insulators and solutions, *Nature*, **423**, 626.
- Van de Walle, C. G., and Neugebauer, J. (2006). Hydrogen in semiconductors, *Annu. Rev. Mater. Sci.*, **36**, 179–201.
- Vanderbilt, D. (1990). Soft self-consistent pseudopotentials in a generalized eigenvalue formalism, *Phys. Rev. B*, **41**, 7892.
- Vaughan, M. P., and Ridley, B. K. (2007a). Effect of non-parabolicity on the density of states for high-field mobility calculations in dilute nitrides, *Phys. Status Solidi C*, **4**, 686.
- Vaughan, M. P., and Ridley, B. K. (2007b). Electron-nitrogen scattering in dilute nitrides, *Phys. Rev. B*, **75**, 195205.
- Vavilov, V. S., Kiv, A. E., and Niyazova, O. (1975). The subthreshold radiation effects in semiconductors, *Phys. Status Solidi A*, **32**, 11–33.
- Vegard, L. (1921). Die konstitution der mischkristalle und die raumfüllung der atome, *Z. Phys.*, **5**, 17.
- Virkkala, V., Havu, V., Tuomisto, F., and Puska, M. J. (2012). Hybrid functional study of band structures of  $\text{GaAs}_{1-x}\text{N}_x$  and  $\text{GaSb}_{1-x}\text{N}_x$  alloys, *Phys. Rev. B*, **85**, 085134.
- Vogl, P. (1984). Predictions of deep impurity level energies in semiconductors, *Adv. Electron. Electron Phys.*, **62**, 101.
- Volz, K., Koch, J., Kunert, B., and Stoltz, W. (2003). Doping behaviour of Si, Te, Zn and Mg in lattice-matched (GaIn)(NAs)/GaAs bulk films, *J. Cryst. Growth*, **248**, 451.
- Volz, K., Torunski, T., Lackner, D., Rubel, O., Stoltz, W., Baur, C., Mueller, S., Dimroth, F., and Bett, A. W. (2007). Material development for improved 1 eV (GaIn)(NAs) solar cell structures, *J. Sol. Energy Eng.*, **129**, 266.
- Vorona, I. P., McHedlidze, T., Dagnelund, D., Buyanova, I. A., Chen, W. M. and Kohler, K. (2006). Optically detected magnetic resonance studies of point defects in Ga(Al)NAs, *Phys. Rev. B*, **73**, 125204.
- Vouilloz, F., Oberli, D. Y., Dupertuis, M. A., Gustafsson, A., Reinhardt, F. and Kapon, E. (1997). Polarization anisotropy and valence band mixing in semiconductor quantum wires, *Phys. Rev. Lett.*, **78**, 1580.
- Wang, X. J., Buyanova, I. A., Zhao, F., Lagarde, D., Balocchi, A., Marie, X., Tu, C. W., Harmand, J. C., and Chen, W. M. (2009a). Room-temperature defect-engineered spin filter based on a non-magnetic semiconductor, *Nat. Mater.*, **8**, 198.
- Wang, X. J., Puttisong, Y., Tu, C. W., Ptak, A. J., Kalevich, V. K., Egorov, A. Y., Geelhaar, L., Riechert, H., Chen, W. M., and Buyanova, I. A. (2009b).

- Dominant recombination centers in Ga(In)NAs alloys: Ga interstitials, *Appl. Phys. Lett.*, **95**, 241904.
- Warren, B. E. (1990). *X-ray Diffraction* (Dover, New York).
- Wei, S. H., and Zunger, A. (1996). Giant and composition-dependent optical bowing coefficient in GaAsN alloys, *Phys. Rev. Lett.*, **76**, 664.
- Weinstein, B. A., Stambach, S. R., Ritter, T. M., Maclean, J. O., and Wallis, D. J. (2003). Evidence for selective delocalization of N-pair states in dilute  $\text{GaAs}_{1-x}\text{N}_x$ , *Phys. Rev. B*, **68**, 035336.
- Weiringen, A. V., and Warmholtz, N. (1956). On the permeation of hydrogen and helium in single crystal silicon and germanium at elevated temperatures, *Physica*, **22**, 849–865.
- Wen, L., Bekisli, F., Stavola, M., Fowler, W. B., Trotta, R., Polimeni, A., Capizzi, M., Rubini, S., and Martelli, F. (2010). Detailed structure of the H-N-H center in  $\text{GaAs}_{1-y}\text{N}_y$  revealed by vibrational spectroscopy under uniaxial stress, *Phys. Rev. B*, **81**, 233201.
- Wen, L., Stavola, M., Fowler, W. B., Trotta, R., Polimeni, A., Capizzi, M., Bisognin, G., Berti, M., Rubini, S., and Martelli, F. (2012). Microscopic origin of compressive strain in hydrogen-irradiated dilute  $\text{GaAs}_{1-y}\text{N}_y$  alloys: role of N-H<sub>n</sub> centers with n>2 and their thermal stability, *Phys. Rev. B*, **86**, 085206.
- Woicik, J. C., Pellegrino, J. G., Steiner, B., Miyano, K. E., Bompadre, S. G., Sorensen, L. B., Lee, T. L., and Khalid, S. (1997). Bond-length distortions in strained semiconductor alloys, *Phys. Rev. Lett.*, **79**, 5026.
- Wolford, D. J., Bradley, J. A., Fry, K., and Thompson, J. (1984). The nitrogen isoelectronic trap in GaAs, in *Proceedings of the 17<sup>th</sup> International Conference on the Physics of Semiconductors* (Springer, New York), 627.
- Wormington, M., Panaccione, C., Matney, K. M., and Bowen, K. (1999). Characterisation of structures from x-ray scattering data using genetic algorithms, *Philos. Trans. R. Soc. London, A*, **357**, 2827.
- Xin, H. P., and Tu, C. W. (2000). Photoluminescence properties of GaNP/GaP multiple quantum wells grown by gas source molecular beam epitaxy, *Appl. Phys. Lett.*, **77**, 2180.
- Xin, H. P., Tu, C. W., and Geva, M. (1999). Annealing behavior of p-type  $\text{Ga}_{0.892}\text{In}_{0.108}\text{N}_x\text{As}_{1-x}$  ( $0 \leq x \leq 0.024$ ) grown by gas-source molecular beam epitaxy, *Appl. Phys. Lett.*, **75**, 1416.
- Yang, X., Heroux, J. B., Mei, L. F., and Wang, W. I. (2001). InGaAsNSb/GaAs quantum wells for 1.55  $\mu\text{m}$  lasers grown by molecular-beam epitaxy, *Appl. Phys. Lett.*, **78**, 4068.

- Yang, X., Jurlovic, M. J., Heroux, J. B., and Wang, W. I. (1999). Molecular beam epitaxial growth of InGaAsN:Sb/GaAs quantum wells for long-wavelength semiconductor lasers, *Appl. Phys. Lett.*, **75**, 178.
- Yoshie, T., Scherer, A., Hendrickson, J., Khitrova, G., Gibbs, H. M., Rupper, G., Ell, C., Shchekin, O. B., and Deppe, D. G. (2004). Vacuum Rabi splitting with a single quantum dot in a photonic crystal nanocavity, *Nature*, **432**, 200.
- Yuan, J., Wang, H., Veldhoven, R. P. J. V., Wang, J., Vries, T. D., Smalbrugge, B., Jin, C. Y., Nouwens, P., Geluk, E. J., Silov, A. Y., and Noetzel, R. (2011). Controlling polarization anisotropy of site-controlled InAs/InP (100) quantum dots, *Appl. Phys. Lett.*, **98**, 201904.
- Zangrando, M., Finazzi, M., Zacchigna, M., Cocco, D., Rochow, R. and Parmigiani, F. (2004). Polarized high-brilliance and high-resolution soft x-ray source at ELETTRA: the performance of beamline BACH, *Rev. Sci. Instrum.*, **75**, 31.
- Zavada, J. M., Jenkinson, H. A., Sarkis, R. G., and Wilson, R. G. (1985). Hydrogen depth profiles and optical characterization of annealed, proton-implanted n-type GaAs, *J. Appl. Phys.*, **58**, 3731.
- Zhang, Y., Fluegel, B., Mascarenhas, A., Xin, H. P., and Tu, C. W. (2000). Optical transitions in the isoelectronically doped semiconductor GaP:N: an evolution from isolated centers, pairs, and clusters to an impurity band, *Phys. Rev. B*, **62**, 4493.
- Zhang, Y., and Mascarenhas, A. (2000). Isoelectronic impurity states in GaAs:N, *Phys. Rev. B*, **61**, 15562.
- Zory, P. S. J. (1993). *Quantum Well Lasers* (Academic Press, New York).
- Zwiller, V., Pistol, M., Hessman, D., Cederstrom, R., Seifert, W. and Samuelson, L. (1999). Time-resolved studies of single semiconductor quantum dots, *Phys. Rev. B*, **59**, 5021.

# Index

- absorption 78, 161, 173, 177–179, 181–183, 186  
infrared 39, 225  
intervalence band 24  
nitrogen mode 133  
phonon-related 132  
photoelectric 168, 169, 173  
photon 73, 255  
absorption edge 39, 168, 170, 173, 177–179  
acceptors 76, 100, 103, 110–112, 121  
compensated 121  
ionized 111, 112  
negative 121  
shallow 111, 112, 116  
activation energy 43–45, 47, 67, 73, 95–97, 126, 234, 235  
alloys 1–5, 8, 10, 16, 28, 29, 32, 37, 52, 54, 56, 85–87, 90, 100–103, 130, 132  
bulk 198  
quaternary 19  
semiconductor 1, 5, 15, 16  
alloy scattering 4, 15  
annealing 47, 85, 92, 122, 126, 155, 156, 195, 207, 209, 212, 213, 224, 225, 257  
aggressive 224  
laser 255  
moderate 116  
approximation 173, 233  
Born wave 15  
generalized gradient 105  
gradient-corrected 139  
local density 102  
muffin-tin 173, 176  
virtual crystal 101  
atomic arrangement 111, 125, 126  
atomic configurations 111, 112, 115, 185  
atomic distances 102, 115, 116  
atomic orbitals 168, 202, 203  
atomic species 110, 171, 177, 185, 201, 216  
atoms 16, 17, 31, 32, 65–67, 90–92, 101–104, 107–127, 156–161, 168–177, 185–191, 193, 194, 210, 211, 219, 220, 234, 235, 237–240, 242, 243  
absorbing 172, 176, 179  
anomalous 179, 180  
artificial 229  
defective 185  
excited 172, 185, 216  
free 173  
host 130, 139  
host-type 211
- BAC model *see* band-anticrossing model  
band-anticrossing model (BAC model) 1, 4–10, 14, 15, 28, 29, 56, 113, 114  
band edges 14, 16, 89, 98, 114

- band gap 46, 49, 56, 57, 89, 98, 101, 103, 129–131, 152, 160, 167, 234, 239, 245, 256  
 experimental 103  
 fundamental 256  
 optical 166  
 band-gap bowing 3, 4, 29, 113, 114  
 band-gap energy 33, 34, 38, 54, 65, 73, 76, 85, 89, 98, 188, 208, 229, 230  
 band-gap engineering (BGE) 230, 238, 239, 241, 243  
 band-gap shifts 130, 133, 136, 152, 159  
 barriers 141, 142, 146, 148, 216, 241, 243, 247  
 chemical reaction 105  
 finite-height 244  
 hydrogenated 243, 247, 248  
 impenetrable 237  
 reorientation 148  
 beam 216–219, 230, 253, 254, 259  
 collimated 214  
 deutron 216, 217  
 laser 48–50, 52  
 monochromatic 213  
 photon 174  
 BGE *see* band-gap engineering  
 bonds 47, 109, 127, 139, 146, 188, 199, 200, 253  
 chemical 52  
 dangling 32, 116, 191–193  
 dopant–neighbor 127  
 Bragg peak 178, 213, 214, 220, 221, 224  
 carrier density 17, 21, 23, 25  
 carrier recombination 41, 62–64, 80, 89, 90, 98, 164, 254  
 carriers 16, 25, 54, 55, 61, 240, 243, 253  
 CB *see* conduction band  
 CB dispersion 7, 8, 21  
 CBE *see* conduction band edge  
 CBE energy 6, 16, 18  
 CB minimum 7, 54, 56, 57, 60, 61, 65, 68, 71, 72  
 CCD *see* charge-coupled device  
 charge-coupled device (CCD) 35, 36, 184  
 charge states 79, 90, 103, 109, 116, 117, 119–121, 139, 141, 191–194  
 clusters 4, 11, 12, 32, 43, 45, 57, 59–62, 65–67, 74, 188, 201, 262  
 cluster states 12, 15, 18, 64, 65  
 complexes 10, 31, 32, 50, 53, 90, 91, 97–102, 116, 117, 119–127, 131, 188–194, 209, 210, 224, 225, 229, 234, 237  
 charge transfer 105  
 dihydrogen 189, 254  
 hydrogen–impurity 76  
 hydrogen–nitrogen 116  
 interstitial 95  
 monohydrogen 189  
 conduction band (CB) 5, 21, 22, 33, 39, 61, 68, 71, 89, 100, 113, 114, 201–203, 242, 249  
 conduction band edge (CBE) 2–4, 7, 10–12, 14, 15, 18, 21, 29, 92, 97, 98  
 core hole 168, 172, 177, 182, 201, 202  
 Coulomb barrier 215, 216  
 DAFS *see* diffraction anomalous fine structure  
 dangling bond (DB) 32, 76, 111, 116, 117, 127, 191–193, 199  
 DB *see* dangling bond  
 Debye–Waller factor 200  
 decay 79, 80, 168, 172, 216, 237, 241

- radioactive 217
- single exponential 224
- deep-level transient spectroscopy
  - (DLTS) 77, 95, 96, 98
- defects 27, 32, 75–82, 84–87, 90–96, 98, 100, 102, 103, 129–131, 135–142, 144–147, 157–160, 163, 190, 191, 219
  - active 262
  - complex 158
  - interstitial-related 81
  - intrinsic 77
  - isolated 103
  - paramagnetic 78, 79
  - radiative 228
  - trigonal 131
- density 14, 21–23, 25–27, 35, 42, 60, 61, 65, 69, 102, 104, 105, 108, 132, 157, 161, 167
  - areal 217
  - electron charge 65
  - electronic 105
  - excitation power 64
  - finite 9
  - hole 25
  - infinite 9
- density functional theory (DFT)
  - 102, 104–106, 139, 157, 161, 167, 187, 194
- DFT *see* density functional theory
- diffraction anomalous fine structure (DAFS) 162, 168, 177, 178, 180, 195, 197, 198, 204
- diffusion 51, 52, 91, 227, 232–234, 237, 239
  - carrier 242
  - controlled 242
  - thermal 35
- diffusion coefficients 232, 233
- distribution 3, 4, 10–12, 14, 15, 17, 47, 83, 102, 157, 238, 242, 248
  - bond length 201
  - charge density 116
- electromagnetic field 228
- statistical 11, 57
- DLTS *see* deep-level transient spectroscopy
- dopants 32, 111, 113, 172, 176, 185, 213
  - neutral 76
  - shallow 99, 112, 127
- effective mass 3, 5, 7, 12, 13, 16, 18, 21, 22, 33–35, 53–56, 58–62, 65–67, 71–73, 76, 101, 251
- effects
  - band-filling 241
  - damping 201
  - dead-time 172
  - electronic 119, 124–126
  - hydrogen-induced
    - neutralization 49
  - hydrogen-related 187
  - optical 167, 207, 208
  - photonic 48
  - polarization 246
  - quantum confinement 94
- elastic energy 66, 67
- electron gyromagnetic factor 31, 53, 70–72
- electronic properties 1, 3, 29, 31, 32, 34–38, 40, 42, 44, 46, 48, 50, 52–54, 56–72, 74, 161–163
- electronic states 2, 72, 103, 184, 201
- electronic structure 1, 3, 8, 10, 20, 21, 28, 29, 123, 125, 127, 176, 182, 201, 203
- electron paramagnetic resonance (EPR) 79
- electron wavefunction 79, 81, 169
- emission 20, 27, 42, 85, 94, 172, 182, 183, 216, 241, 243, 246

- dot 242, 244
- laser 21
- longer-wavelength 20, 28
- near-band edge 92
- near-band-edge 92, 93
- optical 2
- single-photon 262
- spontaneous 25
- emission energy 45
- energy 5, 6, 8, 9, 11, 12, 14, 15, 37, 38, 40, 41, 48, 59, 60, 65–67, 102–106, 168, 182–184, 186, 187, 215–218, 252–254
- biexciton binding 229
- binding 105, 157
- cluster state 12
- exciton 57
- gap 231
- ground-state 147, 243
- laser photon 52
- peak 42, 49, 50
- photoelectron 170
- quasi-Fermi 22
- resonant-level 11
- resonant-state 4
- threshold displacement 89
- zero-point 149
- energy barriers 91, 148, 209, 210, 224, 225
- energy gap 2, 4, 6, 7, 12, 19, 20, 28, 33, 39, 100, 101, 103, 110, 111, 123, 125, 253, 256
- epilayers 37–40, 42, 44, 48, 67, 69, 83, 89, 186, 193, 199, 200, 220, 221, 224, 261
- deuterated/hydrogenated 192
- hydrogenated 200
- passivated 224
- untreated 187, 200
- EPR *see* electron paramagnetic resonance
- EXAFS *see* extended X-ray absorption fine structure
- EXAFS function 174, 175
- EXAFS region 173–175
- EXAFS signal 176, 199
- EXAFS spectrum 174, 175
- experiments 78, 79, 81, 83, 130, 131, 139, 140, 146, 147, 149, 150, 152, 159, 160, 162, 180–182, 190, 193, 196, 198
- laser dissociation 73
- laser-writing 50
- optical spectroscopy 31
- synchrotron radiation 205
- extended X-ray absorption fine structure (EXAFS) 170, 173, 176, 195, 197
- Fermi energy 103, 109, 119
- Fermi level 102, 131
- Fick's equation 233, 234
- fluorescence yield (FY) 171, 172
- FMS *see* full multiple-scattering
- FMS simulations 187, 193, 194
- Fourier transform 130, 175, 200
- frequencies 43, 134, 148, 150, 154, 156, 158, 178, 235
- angular 184
- mode 135
- phonon 148
- scissoring 123, 124
- vibrational 122, 141, 157
- full multiple-scattering (FMS) 176
- FY *see* fluorescence yield
- Gaussian model 175
- growth 19, 27, 32, 83, 261
- epitaxial 191, 210, 229, 261
- exponential 19
- pseudomorphic 193
- high-resolution X-ray diffraction (HRXRD) 115, 122, 152, 153, 157, 167, 191, 194, 207–211, 213, 218, 221, 222

- H-impurity complexes 111  
 H-induced effects 32, 232  
 H-induced recovery 85, 256, 258  
 H-induced variation 101, 258  
*HRXRD* *see* high-resolution X-ray diffraction  
 hydrogenated samples 14, 35, 44, 54–56, 63–65, 70, 73, 83, 92, 134, 164, 190, 197–199, 253, 256  
 hydrogenation 3, 31, 32, 34–36, 38–40, 42–44, 46–48, 72–74, 83, 85, 89, 90, 95, 97, 161–163, 167, 230, 231  
 postgrowth 75, 82, 85, 86, 94, 96, 97  
 selective 227, 237, 239, 249, 258, 262  
 hydrogen diffusion 232, 233, 238  
 hydrogen doses 38, 40, 54, 62, 63, 83, 164, 195, 239  
 hydrogen effects 42, 43, 45, 47, 49, 51–53, 227  
 hydrogen incorporation 31, 32, 34, 49, 53, 54, 73, 95, 96  
 hydrogen irradiation 132, 136, 152, 167, 186, 204, 208, 219, 230, 252, 254–256, 258  
 hydrogen passivation 65, 92  
 hydrostatic pressures 6, 7, 31, 33, 57–59, 64, 65, 74
- IBA* *see* ion beam analysis  
 impinging photon 168, 182, 183  
 impurities 15, 16, 32, 75, 76, 99, 100, 110, 113, 117, 125, 127, 162, 163, 211, 213, 219, 232, 233, 237, 238  
 amphoteric 100, 110  
 charged 233  
 isoelectronic 99, 101, 117, 127, 162  
 isolated 16
- shallow 100, 101, 110, 111  
 transition metal 106  
 in-plane BGE 231, 238, 239, 241, 252, 254, 259–261, 263  
 intensity 39, 43, 44, 47, 63, 66, 85, 93, 95, 132, 133, 135, 144–146, 155, 171, 172, 244, 246  
 diffracted 178, 180, 181, 183, 196  
 field 70  
 relative 142, 144–147, 151, 190  
 scattered 164  
 spectral-line 156  
 interaction 2, 5, 9, 11, 12, 18, 21, 56, 60, 61, 65, 71, 113, 199, 203, 232, 234  
 hyperfine 81, 82  
 local 125, 127  
 massive 167  
 quantum 33  
 two-level 33  
 ion beam analysis (IBA) 208, 214  
 irradiation 35, 36, 39, 42, 46, 47, 132, 133, 165, 167, 188, 190, 203, 231, 232, 243, 245, 251, 252, 256–259  
 isosurfaces 108, 112, 118  
 isotopes 81, 141, 216, 217
- laser exposure 50, 51  
 laser light 48, 252, 255  
 laser powers 49–52, 73  
 lasers 19, 20, 25, 27, 28, 38, 48, 53  
 long-wavelength 129  
 nitride-based 28  
 tunable 36  
 vertical-cavity 5, 28  
 laser spot 50, 51  
 lattice 39, 66, 73, 126, 152, 153, 160, 166, 229  
 crystalline 219  
 host 33, 211

- perfect 16
- periodic 210
- lattice constant 34, 101, 126, 165, 230
- lattice expansion 241, 247, 249, 256
- lattice overshooting 115, 124, 126
- lattice parameter 115, 163, 166, 167, 178, 190, 198, 199, 210, 211, 213, 220, 221
- layer 42, 79, 155, 156, 196, 207, 209, 214, 220, 221, 223–225, 243, 256
  - absorbing 19
  - active 166, 178
  - hydrogenated 166
  - passivated 222
  - tensile 23
  - tensile-strained 23
- local structure 125, 162, 170, 171, 173, 174, 176, 178, 191, 208
- local vibrational modes (LVMs) 130, 134, 188
- LVMs *see* local vibrational modes
  
- magnetic fields 31, 54, 55, 57, 58, 68–70, 74, 79, 250
- magnetic moments 81
- magnetophotoluminescence 13, 54
- masks 230, 231, 237, 260–262
- method
  - empirical pseudopotential 114
  - experimental 34, 35, 182, 207, 208
  - finite-element 237
  - nanofabrication 250
  - numerical 176
  - spectroscopic 78
- microscope 255
  - optical 53, 255
  - scanning electron 239
- MIR *see* multiple-internal-reflection
- MIR geometry 154–156, 160
- model 2, 6, 8, 9, 12, 13, 15, 71, 72, 97, 101, 112–114, 116, 188–190, 234, 236, 237, 243, 248, 249
  - anticrossing 3, 13
  - atomic-scale 160
  - classical 17
  - complex 199
  - desorption 51
  - oscillator 145
  - pictorial 112
  - scattering 16
  - structural 175
  - theoretical 114, 115, 122
  - two-level 11, 18
- modes 123, 124, 132–137, 141, 172, 173, 262
  - confined cavity 228
  - electron detection 172
  - high-frequency stretch 141
  - out-of-plane 141
  - single-photon counting 36
  - transmission 171
- monochromator 164, 166, 182
  - focal length 35, 36
  - four-crystal 212
- multiple-internal-reflection (MIR) 130
  
- nanophotonics 227–229, 262
- nanostructures 170, 171, 227–229, 237–239, 243, 254, 261, 262
- nearest neighbors (NNs) 196
- neighboring atoms 79, 100, 169, 170, 173–175
- N–H bonds 43, 45, 46, 50, 117, 131, 137, 140, 160, 194, 252, 255

- N–H complexes 29, 31, 39, 42, 43, 50–52, 73, 102–104, 113, 114, 116, 119, 121, 122, 185, 187–189, 194, 195, 198
- N–H defect 40, 190
- nitride alloys 1–5, 10, 14–18, 21, 28, 29, 34, 48, 52, 53, 72, 73, 167, 249
- nitride lasers 4, 19–21, 23–25, 27, 28
- nitride materials 231, 250, 255
- nitrides 31, 32, 34–36, 38–40, 42, 48–50, 66–68, 72–78, 98–100, 159–163, 187, 188, 207–210, 218–222, 224–229, 232–239, 251–253
- nitrogen concentrations 34, 37, 39, 56, 57, 70, 71, 195, 218, 234, 243
- NNs *see* nearest neighbors
- NRA *see* nuclear reaction analysis
- nuclear reaction analysis (NRA) 167, 191, 194, 207–210, 214, 216, 217, 222, 223
- nuclear reactions 209, 216–219, 223
- ODMR *see* optically detected magnetic resonance
- ODMR intensity 86, 95
- ODMR signals 79, 80, 85, 89, 90, 92–95
- ODMR spectra 80, 81, 85–87, 93, 94, 96
- ODMR technique 78–80, 87, 98
- optically detected magnetic resonance (ODMR) 75, 77–81, 85, 90, 92, 98
- optical properties 1, 3, 29, 42, 43, 48, 53, 54, 73, 166, 229, 240
- optical theorem 177, 178
- passivation 40, 42, 66, 67, 76, 99, 101, 102, 104, 111, 112, 122, 123, 125, 127, 224, 225, 252, 255, 256, 258
- electronic 31, 42, 100, 234
- hydrogen-induced 93, 229
- passivation rate 64, 66
- paths 26, 175, 179
- back and forth scattering 174
- free 16, 17, 172, 189
- mean-free 17
- photoelectron-scattering 179
- photoelectron 169, 170, 173, 176, 179, 189
- photoluminescence (PL) 35, 40–42, 44, 47, 70, 74, 79, 97, 115, 163, 186, 191, 195, 201, 225
- photon energy 70, 168, 170, 171, 185
- photons 35, 182, 229, 253, 254
- PL *see* photoluminescence
- PL bands 41, 65, 92, 93, 164
- PL emissions 79, 80, 85, 86, 91, 92
- PL Intensity 37, 43–45, 79, 85, 86, 90, 92, 93, 164, 241
- PL spectra 37–39, 41–45, 47, 57, 58, 62–64, 66, 67, 89, 164, 165, 240, 249
- polarization 79, 143, 146, 175, 179, 181
- polarized light 130, 142, 146, 147
- properties 1–6, 8, 10, 12, 14–16, 18, 20, 31, 74–76, 99, 100, 102, 107–109, 207, 208, 227, 228, 258, 259
- alloy 76
- chemical 107, 113
- electrical 258
- local 116
- magnetic 106
- physical 32, 60, 73, 163, 190, 201, 204, 229, 237
- polarization 245, 248, 249, 256

- semiconductor 100  
structural 126, 191, 207, 208,  
210, 220, 228, 236, 249
- QDs *see* quantum dots  
quantum confinement 231, 241,  
243  
quantum dots (QDs) 228, 229,  
242, 250, 260  
quantum wells (QWs) 2, 23, 24,  
37, 42, 76, 178, 195–197, 199  
QWs *see* quantum wells  
QWs  
hydrogenated 198  
thin 166, 178
- RBS *see* Rutherford back scattering  
spectroscopy  
RCs *see* rocking curves  
recombination 25, 54, 55, 68, 77,  
79, 80, 82, 92, 244  
bound-hole 54  
charged/multiple-exciton 244  
defect-related 25  
free-exciton 240, 241  
radiative 25, 38, 80, 85, 94, 262  
resonance-enhanced 86  
recombination bands 56, 59, 63,  
67  
refractive index 230, 256–258  
regions 16, 34, 41, 42, 68, 70, 110,  
170, 189, 199, 230, 242, 243,  
246, 254, 259  
deuterated 42  
extended energy 179  
high-charge-density 108  
high-density 108  
high-energy 251  
laser-treated 50  
light-emitting 49  
low-charge-density 110  
low-density 108, 112
- nanometer-sized 238  
near-edge 170, 177, 193  
near-surface 172  
subgap 257  
terahertz 34  
wavelength 34
- resonances 52, 81, 89, 255  
electron paramagnetic 79  
resonant defect levels 2, 3, 5, 10,  
29
- resonant states 6, 9
- rocking curves (RCs) 153, 154,  
220–222, 224–226
- Rutherford back scattering  
spectroscopy (RBS) 214, 215,  
217
- sample holder 219, 224  
sample hydrogenation 35, 244  
samples 35–43, 47–52, 54–65,  
68–72, 74, 138, 139, 146, 147,  
150, 151, 153–156, 164–166,  
202, 212, 213, 217–219,  
221–225, 235–240  
deuterated 42, 122, 134, 138,  
149  
deuterium-irradiated 132  
electron-irradiated 253  
epilayer 94  
hydrogenated GaNP 77  
isotropic 174  
lower-dose irradiated 42  
masked 230, 260  
nitride 14, 259  
temperature-sensitive 53  
unpatterned 241, 248  
virgin N-containing 46
- scanning electron microscopy  
(SEM) 49, 51, 52
- scanning near-field optical  
microscope (SNOM) 53, 255
- scattering 4, 15–17, 169, 170, 173,  
176, 219

- band-edge 18
- inelastic 184
- n-type carrier 4, 16
- resonant 15
- scattering angle 215
- scattering paths 176, 179
- scattering theory 168
- SDDs *see* Si drift diodes
- SE *see* spontaneous emission
- secondary ion mass spectrometry (SIMS) 35, 42, 83, 191, 209, 236
- SEM *see* scanning electron microscopy
- semiconductor
  - heterostructures 170, 171, 210
  - semiconductor lasers 19, 25
  - semiconductor materials 32, 79, 95, 227
  - semiconductor matrix 100, 232
  - semiconductors 32, 33, 35, 92, 100–102, 162, 163, 168, 169, 171–173, 175–177, 179, 181, 183, 185, 227, 229, 232, 233
    - alloyed 32
    - conventional 232
    - host 100, 101
    - narrow-gap 32
    - nondegenerate 17
  - Si drift diodes (SDDs) 172
  - signals 87, 175, 182, 185–187, 189, 202, 217
    - experimental 189
    - filtered 196
    - luminescence 36
    - multiline 95
    - shell 200
    - single-line 95
  - signal-to-noise ratio 130, 154, 155, 171, 172, 197, 212
  - SIMS *see* secondary ion mass spectrometry
  - SIMS measurements 36, 74, 83, 236, 263
  - simulated ODMR spectra 86, 87, 91, 93
  - simulations 87, 107, 175, 176, 185, 187, 189–195, 197, 199, 201, 203, 213, 220, 221, 236, 239, 240, 261
    - finite-element 248
    - theoretical 103, 114
  - SNOM *see* scanning near-field optical microscope
  - spectra 38, 40, 64, 67, 68, 70, 92, 94, 95, 143, 149, 151, 155, 193, 202, 203, 217, 219
    - bottommost 244
    - experimental 86, 176, 192, 194, 197
    - lowest 155
    - magneto-PL 54, 57
    - photorelectance 7
    - raw 196
    - simulated 93, 189
    - topmost 243
    - vibrational 136, 141
  - spontaneous emission (SE) 25
  - strain
    - extra-compressive 257
    - local 66
    - random 150, 151
  - strain fields 245, 248
  - strain profiles 213, 220, 221
  - stress
    - biaxial anisotropic 248
    - built-in 248
    - tensile 247
  - stress direction 143, 147
  - substrate 19, 186, 193, 195, 196, 210, 211, 213, 224, 244
    - laser 25
    - patterned 251
    - thick 154, 213
  - supercells 10–12, 66, 102–104, 124, 139, 188

- telecom lasers 5, 19, 20  
TEY *see* total electron yield  
thermal annealing 3, 5, 29, 35, 43,  
  46, 47, 73, 77, 83, 85, 167,  
  186, 209, 210, 224, 258  
thermal dissociation 50, 51, 73  
total electron yield (TEY) 171, 172  
transitions 43, 44, 54, 55, 67–70,  
  78, 79, 142, 144–146, 163,  
  183  
carbon impurity 254  
dipole-allowed 68  
magnetic dipole 81, 82  
radiative 248  
spin-forbidden 81  
spin-resonant 79
- uniaxial stress 130, 142, 152, 160
- valence band maximum (VBM) 33,  
  202–204  
VBM *see* valence band maximum  
VCA *see* virtual crystal  
  approximation  
VCSELs *see* vertical-cavity surface  
  emitting lasers  
vertical-cavity surface emitting  
  lasers (VCSELs) 19, 245  
virtual crystal approximation  
  (VCA) 101
- wavefunctions 11, 16, 55, 61, 168  
  carrier 250  
  electron-hole 241  
  final-state 169, 173  
  many-body 174  
  scattered 169
- wavelengths 2, 19, 27, 37, 51, 169,  
  170, 228  
photon 52  
telecom 252
- XAFS *see* X-ray absorption fine  
  structure  
XAFS analysis 178  
XAFS experiments 196  
XAFS measurements 171  
XANES *see* X-ray absorption  
  near-edge structure  
XANES  
  fluorescence 190  
  fluorescence-mode 185  
  interpretation links 177  
  nitrogen-edge 189  
  simulated 187  
XANES experiments 186, 191,  
  195  
XANES spectra 176, 188, 193,  
  202
- XES *see* X-ray emission  
  spectroscopy
- X-ray absorption fine structure  
  (XAFS) 168, 170–173, 178,  
  182, 184, 185, 198, 204
- X-ray absorption near-edge  
  structure (XANES) 140, 152,  
  153, 157, 170, 173, 176, 185,  
  188–191, 193–195, 201–203,  
  205
- X-ray diffraction (XRD) 101,  
  161–163, 165, 167, 178, 185,  
  186, 195, 198, 199
- X-ray emission spectroscopy  
  (XES) 162, 168, 182, 184,  
  204, 205
- XRD *see* X-ray diffraction